



## PRODUCT SPECIFICATION SHEET

### Actidione Agar w/ Actidione (DM006)

#### Intended Use

Actidione Agar w/ Actidione (DM006) is recommended for the enumeration of bacteria in specimens containing large numbers of yeasts and moulds.

#### Product Summary and Explanation

Actidione Agar was formulated by Green and Gray,<sup>(1)</sup> which may be used for microbiological investigation during brewing and baking. Actidione (Cycloheximide) at a concentration of 0.001% permits the growth of bacteria but inhibits the growth of most yeasts and moulds except dermatophytes. Media containing this antibiotic are invaluable for the enumeration and detection of bacteria in specimens containing large numbers of yeasts and moulds. This medium may be used for the estimation of bacterial contamination of pitching yeast. Actidione Agar with added penicillin and streptomycin is also valuable as a selective medium for the isolation of dermatophytes. Green and Gray employed their medium at two different reactions, pH 5.5 and pH 6.5, the latter may be attained by adding approximately 16ml of sterile 1.5% sodium carbonate to each litre of molten medium.

#### Principles of the Procedure

Actidione Agar contains casein enzymic hydrolysate which provides nitrogen, carbon, and amino acids essential growth nutrients. Yeast extract is a rich source of vitamin B complex. Dextrose in high amount along with mineral salts at acidic pH favour sugar fermentation. Monopotassium phosphates buffer the medium. Potassium Chloride, calcium chloride, and ferric chloride are essential ions and help to maintain osmotic balance. Magnesium Sulfate provides divalent cations and sulfate. Bromo cresol green acts as an indicator dye. Actidione permits the growth of bacteria but inhibits the growth of most yeasts and moulds except dermatophytes.

#### Formula / Liter

Ingredients	Gms / Liter
Casein enzymic hydrolysate	5.00
Yeast extract	4.00
Dextrose	50.00
Monopotassium phosphate	0.55
Potassium chloride	0.425
Calcium chloride	0.125
Magnesium sulphate	0.125
Ferric chloride	0.0025
Magnesium sulphate	0.0025
Bromo cresol green	0.022
Actidione (Cycloheximide)	0.01
Agar	15.00
Final pH: 5.5 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

#### Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.
3. Actidione (Cycloheximide) is very toxic. Avoid skin contact or aerosol formation and inhalation.





## PRODUCT SPECIFICATION SHEET

### Directions

1. Suspend 75.26 grams of the medium in one liter of distilled water.
2. Heat to boiling to dissolve the medium completely.
3. Autoclave at 121°C, 15 psi pressure, for 15 minutes / validated cycle.
4. Cool to 45°C. Mix well before pouring into sterile Petri plates.

### Quality Control Specifications

<b>Dehydrated Appearance</b>	Light yellow to light green homogeneous free flowing powder
<b>Prepared Medium</b>	Greenish blue clear to slightly opalescent gel forms in Petri plates
<b>Reaction of 7.53% solution</b>	pH 5.5 ± 0.2 at 25°C
<b>Gel Strength</b>	Firm, comparable with 1.5% Agar gel

**Expected Cultural Response:** Cultural characteristics observed after an incubation at 30°C for 40-48 hours.

Sr. No.	Organisms	Results to be achieved		
		Inoculum (CFU)	Growth	Recovery
1.	<i>Escherichia coli ATCC 25922</i>	50-100	good-luxuriant	≥50%
2.	<i>Lactobacillus fermentum ATCC 9338</i>	50-100	good-luxuriant	≥50%
3.	<i>Proteus mirabilis ATCC 25933</i>	50-100	good-luxuriant	≥50%
4.	<i>Saccharomyces cerevisiae ATCC 9763</i>	≥10 <sup>3</sup>	inhibited	0%
5.	<i>Saccharomyces uvarum ATCC 28098</i>	≥10 <sup>3</sup>	inhibited	0%

The organisms listed are the minimum that should be used for quality control testing.

### Test Procedure

Refer to appropriate references for standard test procedures.

### Results

Refer to appropriate references and standard test procedures for interpretation of results.

### Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

### Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

### Limitations of the Procedure

1. For identification, organisms must be in pure culture. Morphological, biochemical and/or serological tests should be performed for final identification.
2. Consult appropriate texts for detailed information and recommended procedures.

### Packaging

Product Name : Actidione Agar w/ Actidione

Product Code : DM006

Available Pack sizes : 500gm





## PRODUCT SPECIFICATION SHEET

### References

1. Green, S.R. and Gray, P.P. 1950, Wallerstein Lab. Communication 13,357.

### Further Information




For further information please contact your local MICROMASTER Representative.



### MICROMASTER LABORATORIES PRIVATE LIMITED

Unit 38/39, Kalpataru Industrial Estate,  
Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.  
Ph: +91-22-25895505, 4760, 4681. Cell: 9320126789.  
Email: [micromaster@micromasterlab.com](mailto:micromaster@micromasterlab.com)

QAD/FR/024, Rev.00/01.01.2018

Prepared By	Checked By	Approved By
 01.01.2018	 01.01.2018	 01.01.2018
Microbiologist	Head Quality Control	Head Quality Assurance

### Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

