



# PRODUCT SPECIFICATION SHEET

## Blood Agar Base /Infusion Agar (DM040)

### Intended Use

Blood Agar Base / Infusion Agar (DM040) is used with blood for the isolation and cultivation of a wide variety of microorganisms.

### Product Summary and Explanation

Blood Agar Bases are typically supplemented with 5-10% sheep, rabbit, or horse blood, and used for isolating, cultivating, and determining hemolytic reactions of fastidious pathogenic microorganisms. Without enrichment, this blood agar base can be used as a general purpose medium. Blood Agar Base is a nutritious base, and a variety of supplements, including blood, are often added to enhance this medium. Blood Agar Base is formulated to achieve good growth and improve hemolytic reactions of important pathogenic bacteria.

### Principles of the Procedure

Blood Agar Base is prepared using specially selected raw materials to support good growth of a wide variety of fastidious microorganisms. Blood Agar Base contains Brain heart infusion and Tryptose, which provide nitrogen, carbon, and essential vitamins to stimulate organism growth. Along with essential nutritive properties, the peptones improve and enhance hemolysin production. Sodium Chloride maintains the osmotic balance of the medium. Agar is the solidifying agent. In general, blood agar bases are relatively free of reducing sugars, which have been reported to adversely influence the hemolytic reactions of  $\alpha$ -hemolytic streptococci.<sup>(3)</sup> Supplementation with blood (5 - 10%) provides additional growth factors for fastidious microorganisms, and aids in determining hemolytic reactions. Hemolytic patterns may vary with the source of animal blood and the type of basal medium used.<sup>(4)</sup>

### Formula / Liter

Ingredients	Gms / Litre
Beef heart infusion	10.00
Tryptose	10.00
Sodium chloride	5.00
Agar	15.00
Final pH: 7.3 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

### Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.

### Directions

1. Suspend 40 g of the medium in one liter of distilled water.
2. Heat to boiling to completely dissolve the medium.
3. Autoclave at 121°C, 15 psi pressure, for 15 minutes.
4. Prepare 5% blood agar by aseptically adding the appropriate volume of sterile defibrinated blood to melted sterile molten agar base, cooled to 45 - 50°C.

### Quality Control Specifications

Dehydrated Appearance	Cream to yellow colored, homogeneous, free flowing powder
-----------------------	---



## PRODUCT SPECIFICATION SHEET

<b>Solution</b>	4% Solution in Distilled or deionized water is soluble on boiling, Light to Amber colored, and very slightly to slightly opalescent.
<b>Prepared Medium</b>	Without Blood - light to medium amber - slightly opalescent. With 5% sheep blood the medium is cherry red and opaque.
<b>Reaction of 4.0% Solution</b>	pH 7.3 ± 0.2 at 25°C
<b>Gel Strength</b>	Firm, compared to 1.5% Agar Gel.

### Expected Cultural Response:

Sr. No.	Organisms	Results to be achieved		
		Growth w/o blood	Growth w/ blood	Haemolysis
1.	<i>Escherichia coli</i> ATCC 25922	good-luxuriant	good-luxuriant	slight beta hemolysis
2.	<i>Neisseria meningitidis</i> ATCC 13090	fair	good-luxuriant	none
3.	<i>Staphylococcus aureus</i> ATCC 25923	good-luxuriant	good-luxuriant	Beta haemolysis
4.	<i>Staphylococcus epidermidis</i> ATCC 12228	good-luxuriant	good-luxuriant	none
5.	<i>Streptococcus pneumonia</i> ATCC 6303	fair to good	good-luxuriant	Alpha haemolysis
6.	<i>Streptococcus pyogenes</i> ATCC 19615	fair to good	good-luxuriant	Beta haemolysis

The organisms listed are the minimum that should be used for quality control testing.



**Beta Hemolysis**

**Alpha Hemolysis**

**Gamma Hemolysis**

### Test Procedure

- Inoculate directly onto the surface of the medium. Streak for isolation with an inoculating loop, and stab the agar several times to deposit beta-hemolytic streptococci beneath the agar surface. Subsurface growth will display the most reliable hemolytic reactions.<sup>(4)</sup>
- Incubate plates aerobically, anaerobically, or under conditions of increased CO<sub>2</sub> (5 - 10%).

### Results

Examine the medium for growth and hemolytic reactions after 18 - 24 and 48 hours incubation. There are four types of hemolysis on blood agar media described as:<sup>(4)</sup>

- Alpha hemolysis (α) is the reduction of hemoglobin to methemoglobin in the medium surrounding the colony. This produces a green discoloration of the medium.
- Beta hemolysis (β) is the lysis of red blood cells, producing a clear zone surrounding the colony.
- Gamma hemolysis (γ) indicates no hemolysis.
- Alpha-prime-hemolysis (Û) is a small zone of complete hemolysis, surrounded by an area of partial lysis.

### Storage



## PRODUCT SPECIFICATION SHEET

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

### Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

### Limitations of the Procedure

1. The incubation environment can influence hemolytic reactions of beta-hemolytic streptococci.<sup>(4)</sup>
2. Hemolytic reactions of some strains of group D streptococci have been shown to be affected by differences in animal blood.<sup>(4)</sup>

### Packaging

**Product Name : Blood Agar Base**

**Product Code : DMO40**

**Available Pack sizes : 100gm / 500gm**

### References

1. United States Pharmacopeial Convention. 1995. The US pharmacopeia, 23rd ed. The US Pharmacopeial Convention, Rockville, MD.
2. Vanderzant, C., and D. F. Splittstoesser (eds.). 1992. Compendium of methods for the microbiological examination of foods, 3rd ed. American Public Health Association, Washington, D.C.
3. Casman, E. P. 1947. A noninfusion blood agar for neisseriae, pneumococci and streptococci. *Am. J. Clin.* 17:281-289.
4. Ruoff, K. L. 1995. *Streptococcus*, p. 299-305. In P. R. Murray, E. J. Baron, M. A. Pfaller, F. C. Tenover, and R. H. Tenover (eds.). *Manual of clinical microbiology*, 6th ed. American Society for Microbiology, Washington, D.C.
5. Isenberg, H. D. (ed.). 1992. Interpretation of aerobic bacterial growth on primary culture media, *Clinical microbiology procedures handbook*, vol. 1 p. 1.61-1.67. American Society for Microbiology, Washington, D.C.

### Further Information

For further information please contact your local MICROMASTER Representative.



#### MICROMASTER LABORATORIES PRIVATE LIMITED

Unit 38/39, Kalpataru Industrial Estate,  
Off G. B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.  
Ph: +91-9320126789/9833630009/9819991103  
Email: [sales@micromasterlab.com](mailto:sales@micromasterlab.com)

DMO40PSS, QAD/FR/024, Rev.00

#### Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

