



PRODUCT SPECIFICATION SHEET

Dextrose Peptone Broth (DM083)

Intended Use

Dextrose Peptone Broth (DM083) is recommended for routine sterility testing, cultivation of fastidious organisms and enumeration of thermophiles from canned foods.

Product Summary and Explanation

Peptone Water is a minimal growth medium, particularly suitable as a substrate in the study of indole production. Peptic digest of animal tissue used in Peptone Water is rich in tryptophan content. Peptone Water is also utilized as a base for studying patterns of carbohydrate fermentation with the addition of sugar and indicators such as bromocresol purple, phenol red or bromothymol blue. Peptone Water with Phenol Red is recommended^(1, 2, 3) for studying the ability of an organism to ferment a specific carbohydrate which aid in differentiation of genera and species. The formulation of Peptone Water makes it useful for cultivating non-fastidious organisms.⁽¹⁾ This medium is recommended to study fermentation reactions of *Yersinia enterocolitica*. *Yersinia enterocolitica* is a gram negative bacterium, belonging to the family *Enterobacteriaceae*. This organism is the causative agent of a zoonotic disease yersiniosis occurring in humans as well as wide array of animals such as cattle, pigs and birds.⁽⁴⁾ Peptone Water with pH adjusted to 8.4 is suitable for the cultivation and enrichment of *Vibrio* species.

Principles of the Procedure

Dextrose Peptone Broth contains peptic digest of animal tissue which provides essential nutrients necessary for growth. Sodium chloride maintains the osmotic balance of the medium. Phenol red is added as a pH indicator. Fermentation ability of microorganisms is studied by addition of carbohydrates separately to the basal medium before or after sterilization, such as saccharose, rhamnose, salicin, glucose, dextrose etc. at a concentration of 0.5%. Most of the end products of carbohydrate fermentation are organic acids, which, in the presence of phenol red, show a colour change of the medium from red to yellow. If desired, Durhams tube may be used to detect the gas production if produced. The addition of some sugars can lower the pH of the medium. That can be adjusted with sterile 0.1 N NaOH.

Formula / Liter

Ingredients	Gms / Liter
Peptic digest of animal tissue	20.00
Sodium chloride	5.00
Dextrose	10.00
Final pH: 7.2 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.

Directions

1. Suspend 35 grams of the medium in one liter of distilled water.
2. Heat if necessary to dissolve the medium completely.
3. Autoclave at 121°C, 15 psi pressure, for 15 minutes / validated cycle.





PRODUCT SPECIFICATION SHEET

Quality Control Specifications

Dehydrated Appearance	Cream to light yellow homogeneous free flowing powder
Prepared Medium	Light yellow coloured, clear to slightly opalescent solution in tubes
Reaction of 3.5% Solution	pH : 7.2 ± 0.2 at 25°C
Gel Strength	Not Applicable

Expected Cultural Response: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Sr. No.	Organisms	Results to be achieved	
		Inoculum (CFU)	Recovery
1.	<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant
2.	<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	good-luxuriant
3.	<i>Staphylococcus aureus</i> ATCC 25923	50-100	good-luxuriant
4.	<i>Streptococcus pyogenes</i> ATCC 19615	50-100	good-luxuriant

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Refer appropriate references for standard test procedures.

Results

Refer appropriate references and procedures for interpretation of results.

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. For identification, organisms must be in pure culture. Morphological, biochemical and/or serological tests should be performed for final identification.
2. Consult appropriate texts for detailed information and recommended procedures.





PRODUCT SPECIFICATION SHEET

Packaging

Product Name : Dextrose Peptone Broth

SProduct Code : DM083

Available Pack sizes : 500gm

083

1. MacFaddin J. F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.
2. Finegold S. M. and Baron E. J., 1986, Bailey and Scotts Diagnostic Microbiology, 7th Ed., The C.V. Mosby Co., St. Louis.
3. Murray P. R., Baron J. H., Pfaller M. A., Jorgensen J. H. and Tenover F. C., (Eds.), 2003, Manual of Clinical Microbiology, 8th Ed., American Society for Microbiology, Washington, D.C.
4. Collins FM (1996). Pasteurella, and Francisella. In: Barron's Medical Microbiology (Barron S et al., eds.) 4th ed., Univ of Texas Medical Branch.

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITED

DM083SS,QAD/FR/024,Rev.00/01.01.2018

Unit 38/39, Kalpataru Industrial Estate,

Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.

Ph: +91-22-25895505, 4760, 4681. Cell: 9320126789.

Email: micromaster@micromasterlab.com

sales@micromasterlab.com

	Checked By	Approved By
		
Microbiologist	Head Quality Control	Head Quality Assurance

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

