



PRODUCT SPECIFICATION SHEET

Eosin Methylene Blue Agar, Levine (EMB Agar, Levine) (DM094)

Intended Use

EMB Agar, Levine (DM094) is recommended for the isolation, enumeration and differentiation of members of *Enterobacteriaceae*.

Product Summary and Explanation

Levine EMB Agar was developed by Levine^(1,2) and is used for the differentiation of *Escherichia coli* and *Enterobacter aerogenes* and also for the rapid identification of *Candida albicans*. This medium is recommended for the detection, enumeration and differentiation of members of the coliform group by American Public Health Association^(3, 4, 5). Weld^(6, 7) proposed the use of Levine EMB Agar, with added Chlorotetracycline hydrochloride, for the rapid identification of *Candida albicans* in clinical specimens. A positive identification of *Candida albicans* can be made after 24 - 48 hours incubation at 35 - 37°C in 10% carbon dioxide atmosphere, from specimens such as faeces, oral and vaginal secretions and nail or skin scraping etc. However, the typical appearance is variable. Methylene blue and Eosin-Y inhibit gram-positive bacteria to a limited degree. These dyes serve as differential indicators and help to differentiate between lactose-fermenters and non-fermenters in EMB Agar, Levine.

Principles of the Procedure

Peptic digest of animal tissue provides carbon, nitrogen, and other essential growth nutrients in the medium. Lactose and sucrose are the fermentable carbohydrates which are the sources of energy. Eosin-Y and methylene blue serve as differential indicators in response to the fermentation of lactose by microorganisms. Phosphate buffers the medium. The ratio of eosin-Methylene blue is adjusted to approximately 6:1. These indicators also serve to inhibit the growth of most Gram positive organisms to a limited degree. Coliforms produce blue-black colonies due to the taking up of an eosin-methylene blue dye complex by the bacterial cells when the pH drops. *Salmonella* and *Shigella* colonies are colorless or have a transparent amber color. *Escherichia coli* colonies may show a characteristic green metallic sheen due to the rapid fermentation of lactose. Some gram-positive bacteria, such as fecal streptococci, staphylococci and yeasts, will grow on this medium and usually form pinpoint colonies. A number of non-pathogenic, lactose-nonfermenting gram-negative bacteria will grow on this medium and must be distinguished from the pathogenic bacterial strains by additional biochemical tests.

Formula / Liter

Ingredients	Gms / Litre
Peptic digest of animal tissue	10.00
Lactose	10.00
Dipotassium Phosphate	2.00
Eosin Y	0.40
Methylene Blue	0.065
Agar	15.00
Final pH: 7.1 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT, mainly irritating to eyes, respiratory system, and skin. Handle in accordance with good laboratory hygiene and safety practice. Wash hands before breaks and at the end of workday. To protect, use safety glasses and gloves during handling.
3. Use personal protective equipment. Avoid dust formation. Avoid breathing vapors, mist or gas. Ensure adequate ventilation. Avoid breathing dust.
4. Do not let product enter drains.





PRODUCT SPECIFICATION SHEET

5. After use, prepared plates, specimen containers and other contaminated materials must be sterilized by autoclaving before discarding.

Directions

1. Suspend 37.46 g of the medium in one liter of purified water. Mix until suspension is uniform. Heat to dissolve the medium completely.
2. Dispense and sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes. AVOID OVERHEATING.
3. Cool to 45-50°C and shake the medium in order to oxidize the methylene blue (i.e. to restore its blue colour) and to suspend the flocculent precipitate.
4. If EMB Agar is inoculated on the same day, it may be used without autoclave sterilization.
5. Test samples of the finished product for performance using stable, typical control cultures.
6. Store the medium away from light to avoid photooxidation

Quality Control Specifications

Dehydrated Appearance	Light Pink-Purple, homogeneous, free flowing powder
Solution	3.75% Solution in Distilled or deionized water is soluble on boiling, Reddish purple colored, and slightly hazy.
Prepared Medium	Reddish purple coloured, opalescent gel with greenish cast and finely dispersed precipitate forms in Petri plates
Reaction of 3.75% Solution	pH 7.1 ± 0.2 at 25°C
Gel Strength	Firm, compared to 1.5% Agar Gel.

Expected Cultural Response: Cultural response on EMB Agar, Levine observed after incubation at 35 ± 2°C for 18-48 hours.

Sr. No.	Organisms	Results to be achieved			
		Inoculum (CFU)	Growth	Recovery %	Colour of colony
1.	<i>Enterobacter aerogenes</i> ATCC 13048	50-100	good	40-50%	Pink-red without sheen
2.	<i>Escherichia coli</i> ATCC 25922	50-100	luxuriant	≥50%	blue-black with metallic sheen
3.	<i>Escherichia coli</i> NCTC 9002	50-100	luxuriant	≥50%	blue-black with green metallic sheen
4.	<i>Escherichia coli</i> ATCC 8739	50-100	luxuriant	≥50%	blue-black with green metallic sheen
5.	<i>Candida albicans</i> ATCC 10231	50-100	luxuriant (incubated in 10% CO ₂)	≥50%	colourless
6.	<i>Enterococcus faecalis</i> ATCC 29212	50-100	non-poor	≤10%	colourless
7.	<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	luxuriant	≥50%	colourless
8.	<i>Salmonella Typhimurium</i> ATCC 14028	50-100	luxuriant	≥50%	colourless
9.	<i>Saccharomyces cerevisiae</i> ATCC 9763	50-100	non-poor	≤10%	cream
10.	<i>Staphylococcus aureus</i> ATCC 25923	50-100	non-poor	≤10%	colourless
11.	<i>Staphylococcus aureus</i> ATCC 6538	50-100	non-poor	≤10%	colourless
12.	<i>Pseudomonas aeruginosa</i> ATCC 9027	50-100	non-poor	≤10%	colourless



PRODUCT SPECIFICATION SHEET

--	--	--	--	--	--

The organisms listed are the minimum that should be used for quality control testing.


Test Procedure

1. Observe aseptic techniques.
2. Use standard procedures to obtain isolated colonies from specimens.
3. A nonselective medium should also be streaked to increase the chance of recovery when the population of gram-negative organisms is low and to provide an indication of other organisms present in the specimen.
4. Incubate plates, protected from light, at $35 \pm 2^\circ\text{C}$ for 18-24 hours. If negative after 24 hours, reincubate an additional 24 hours.
5. Confirmatory tests should be further carried out for identification of isolated colonies.

Results

After 18 - 24 hours of incubation at $35 \pm 2^\circ\text{C}$, Typical colonial morphology on EMB Agar is as follows:

Escherichia coli..... Large, blue-black, green metallic sheen
Enterobacter/Klebsiella..... Large, mucoid, blue-black
Proteus..... Large, colorless
Salmonella..... Large, colorless to amber
Shigella..... Large, colorless to amber
Pseudomonas..... Irregular, colorless
 Gram-positive bacteria..... No growth to slight growth



E. coli on EMB Agar, Levine (DM094)

Storage

Store the sealed bottle containing the dehydrated medium at $2-30^\circ\text{C}$. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. Due to nutritional variation, some strains may be encountered that grow poorly or fail to grow on this medium.
2. Biochemical and serological tests are performed for complete identification.

Packaging

Product Name : EMB Agar, Levine

Product Code : DM094

Available Pack sizes : 100gm / 500gm



PRODUCT SPECIFICATION SHEET

References

1. Levine M., 1918, J. Infect. Dis., 23:43.
2. Levine M., 1921, Bull. 62, Iowa State College Engr. Exp. Station.
3. Greenberg A. E., Trussell R. R. and Clesceri L. S. (Eds.), 1998, Standard Methods for the Examination of Water and Wastewater, 20th
 1. Ed., APHA, Washington, D.C.
 4. Marshall R. (Ed.), 1992, Standard Methods for the Examination of Dairy Products, 16th ed., APHA Inc., New York.
 5. Downes F. P and Ito K. (Ed.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., American Public Health Association, Washington, D.C.
 6. Weld J. T., 1952, Arch. Dermat. Syph., 66:691.
 7. Weld J. T., 1953, Arch. Dermat. Syph., 67(5):433.
 8. Howard B. J., 1994, Clinical and Pathogenic Microbiology, 2nd Ed., Mosby Year Book, Inc

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITED

DM094PSS,QAD/FR/024,Rev.00

Unit 38/39, Kalpataru Industrial Estate,

Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.

Ph: +91-9320126789/9833630009/9819991103

Email: sales@micromasterlab.com

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

