

PRODUCT SPECIFICATION SHEET

Kligler Iron Agar (DM127)

Intended Use

Kligler Iron Agar (DM127) is used for the differentiation of members of the *Enterobacteriaceae* on the basis of their ability to ferment dextrose and lactose and to liberate sulfides.

Product Summary and Explanation

Kligler Iron Agar is a modification of Kligler's original formula, a combination of the lead acetate medium described by Kligler⁽¹⁾ and Russells Double Sugar Agar⁽²⁾ and is used as a differentiation medium for typhoid, dysentery and allied bacilli.⁽³⁾ Russell described a new double sugar tube medium containing glucose, lactose, and an indicator, for the isolation of typhoid bacilli from urine and feces. Six years later, Kligler developed a simple lead acetate medium for the differentiation of the typhoid-paratyphoid group. Subsequently, Kligler found lead acetate could detect hydrogen sulfide when combined with Russell double sugar medium for the differentiation of typhoid, paratyphoid, and dysentery groups.⁽⁴⁾ Bailey and Lacey substituted phenol red for Andrade indicator previously used as pH indicator.⁽⁵⁾ Kligler Iron Agar differentiates lactose fermenters from the non-fermenters. It differentiates *Salmonella typhi* from other *Salmonellae* and also *Salmonella paratyphi A* from *Salmonella scottmuelleri* and *Salmonella enteritidis*.⁽⁶⁾ Kligler Iron Agar is recommended for differentiation of enteric gram-negative bacilli from clinical specimens^(7, 8) and food samples.⁽⁹⁾

Fermentation of dextrose results in production of acid, which turns the indicator from red to yellow. Since there is little sugar i.e. dextrose, acid production is very limited and therefore a reoxidation of the indicator is produced on the surface of the medium, and the indicator remains red. However, when lactose is fermented, the large amount of acid produced, avoids reoxidation and therefore the entire medium turns yellow.

Principles of the Procedure

Kligler Iron Agar contains peptic digest of animal tissue, beef and yeast extract which provides nitrogen, carbon, and vitamins required for organism growth. Sodium chloride maintains the osmotic balance of the medium. Lactose and glucose (dextrose) are fermentable sugars which enables the differentiation of species of enteric bacilli. Phenol red is the pH indicator, which exhibits a colour change in response to acid produced during the fermentation of sugars. The combination of ferrous sulphate and sodium thiosulphate enables the detection of hydrogen sulfide production, which is evidenced by a black color either throughout the butt, or in a ring formation near the top of the butt. Lactose non-fermenters (e.g., *Salmonella* and *Shigella*) initially produce a yellow slant due to acid produced by the fermentation of the small amount of glucose (dextrose). When glucose (dextrose) supply is exhausted in the aerobic environment of the slant, the reaction reverts to alkaline (red slant) due to oxidation of the acids produced. The reversion does not occur in the anaerobic environment of the butt, which therefore remains acidic (yellow butt). Lactose fermenters produce yellow slants and butts because of lactose fermentation. The high amount of acids thus produced helps to maintain an acidic pH under aerobic conditions. Tubes showing original colour of the medium indicates the fermentation of neither glucose (dextrose) nor lactose. Gas production (aerogenic reaction) is detected as individual bubbles or by splitting or displacement of the agar by the formation of cracks in the butt of the medium. Pure cultures of suspected organisms from plating media such as MacConkey Agar (DM143), Bismuth Sulphite Agar (DM039) or Deoxycholate Citrate Agar (DM577), SS Agar (DM236) etc. are inoculated on Kligler Iron Agar for identification.

Formula / Liter

Ingredients	Gms / Liter
Peptic digest of animal tissue	15.00
Beef extract	3.00
Yeast extract	3.00
Proteose peptone	5.00
Lactose	10.00
Dextrose	1.00
Ferrous sulphate	0.20
Sodium chloride	5.00
Sodium thiosulphate	0.30

PRODUCT SPECIFICATION SHEET

Phenol red	0.024
Agar	15.00
Final pH (at 25°C) 7.4 ± 0.2	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.

Directions

1. Suspend 57.52 grams of dehydrated powder in 1000 ml distilled water.
2. Heat to boiling to dissolve the medium completely.
3. Mix well and distribute into tubes.
4. Autoclave at 121°C, 15 psi pressure, for 15 minutes / validated cycle.
5. Allow the tubes to cool in slanted position to form slopes with about 1 inch butts.
6. Best reactions are obtained on freshly prepared medium.
7. Do not use screw capped tubes or bottles.

Quality Control Specifications

Dehydrated Appearance	Light yellow to pink homogeneous free flowing powder
Prepared Medium	Red coloured, clear to slightly opalescent gel forms in tubes as slants
Reaction of 5.75% solution	pH 7.4 ± 0.2 at 25°C
Gel Strength	Firm, comparable with 1.5% Agar gel

Expected Cultural Response: Cultural characteristics observed after an incubation at 35-37°C for 18-48 hours.

Sr. No.	Organisms	Results to be achieved					
		Inoculum (CFU)	Growth	Slant	Butt	Gas	H ₂ S
1.	<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	acidic reaction, yellowing of the medium	acidic reaction, yellowing of the medium	positive reaction	Negative reaction, no blackening of medium
2.	<i>Enterobacter aerogenes</i> ATCC 13048	50-100	good-luxuriant	acidic reaction, yellowing of the medium	acidic reaction, yellowing of the medium	positive reaction	Negative reaction, no blackening of medium
3.	<i>Klebsiella pneumoniae</i> ATCC 13883	50-100	good-luxuriant	alkaline reaction, red colour of the medium	acidic reaction, yellowing of the medium	positive reaction	Negative reaction, no blackening of medium
4.	<i>Proteus vulgaris</i> ATCC 6380	50-100	good-luxuriant	alkaline reaction, red colour of the medium	acidic reaction, yellowing of the medium	negative reaction	Positive reaction, blackening of medium
5.	<i>Pseudomonas aeruginosa</i> ATCC 27853	50-100	good-luxuriant	alkaline reaction, red colour of the medium	alkaline reaction, red colour of the medium	negative reaction	Negative reaction, no blackening of medium
6.	<i>Salmonella typhi</i> ATCC 6539	50-100	good-luxuriant	alkaline reaction, red colour of the medium	acidic reaction, yellowing of the medium	negative reaction	Positive reaction, blackening of medium

PRODUCT SPECIFICATION SHEET

7.	<i>Salmonella enteritidis</i> ATCC13076	50-100	good-luxuriant	alkaline reaction, red colour of the medium	acidic reaction, yellowing of the medium	positive reaction	Positive reaction, blackening of medium
8.	<i>Shigella flexneri</i> ATCC12022	50-100	good-luxuriant	alkaline reaction, red colour of the medium	acidic reaction, yellowing of the medium	negative reaction	Negative reaction, no blackening of medium

Sr. No.	Organisms	Results to be achieved					
		Inoculum (CFU)	Growth	Slant	Butt	Gas	H ₂ S
9.	<i>Yersinia enterocolitica</i> ATCC 27729	50-100	good-luxuriant	alkaline reaction, red colour of the medium	acidic reaction, yellowing of the medium	variable reaction	Negative reaction, no blackening of medium
10.	<i>Enterobacter cloacae</i> ATCC 13047	50-100	good-luxuriant	acidic reaction, yellowing of the medium	acidic reaction, yellowing of the medium	positive reaction	negative reaction, no blackening of medium
11.	<i>Citrobacter freundii</i> ATCC 8090	50-100	good-luxuriant	acidic reaction, yellowing of the medium	Acidic reaction, yellowing of the medium	positive reaction	positive reaction, blackening of medium
12.	<i>Salmonella Paratyphi A</i> ATCC 9150	50-100	good-luxuriant	alkaline reaction, red colour of the medium	acidic reaction, yellowing of the medium	positive reaction	negative reaction, no blackening of medium
13.	<i>Salmonella Schottmuelleri</i> ATCC 10719	50-100	good-luxuriant	alkaline reaction, red colour of the medium	acidic reaction, yellowing of the medium	positive reaction	positive reaction, blackening of medium

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

1. To inoculate, carefully touch the center of an isolated colony on an enteric plated medium with a cool, sterile needle, stab into the medium in the butt of the tube, and then streak back and forth along the surface of the slant.
2. Several colonies from each primary plate should be studied separately, since mixed infections may occur. Incubate tubes with loosened caps for 18-24 hours at 35 ± 2°C in an aerobic atmosphere.
3. To enhance the alkaline condition in the slant, free exchange of air must be permitted through the use of a loose closure.
4. If the tube is tightly closed, an acid reaction (caused solely by dextrose fermentation) will also involve the slant.

Results

1. An alkaline slant-acid butt (red/yellow) indicates fermentation of dextrose only.
2. An acid slant-acid butt (yellow/yellow) indicates fermentation of dextrose and lactose.
3. An alkaline slant-alkaline butt (red/red) indicates dextrose and lactose did not ferment (non-fermentor).
4. Cracks, splits, or bubbles in the medium indicate gas production.
5. A black precipitate in butt indicates hydrogen sulfide production.

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

PRODUCT SPECIFICATION SHEET

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. Hydrogen sulfide producing organisms may produce a black precipitate to such a degree that the reaction in the butt is completely masked. If hydrogen sulfide is produced, dextrose is fermented even if it is not observed.
2. Further biochemical tests and serological typing must be performed for definite identification and confirmation of organisms.
3. Pure cultures are essential when inoculating Kligler Iron Agar. If inoculated with a mixed culture, irregular observations may occur.
4. Hydrogen sulfide determinations using Kligler Iron Agar should be limited to members of Enterobacteriaceae

Packaging

Product Name : Kligler Iron Agar

Product Code : DM127

Available Pack sizes : 100gm / 500gm

References

1. Russell F. F., 1911, J. Med. Res., 25:217.
2. Kligler I. J., 1917, Am. J. Publ. Health, 7:1041.
3. Kligler I. J., 1918, J. Exp. Med., 28:319.
4. Kligler, I. J. 1918. Modifications of culture media used in the isolation and differentiation of typhoid, dysentery, and allied bacilli. J. Exp. Med. 28:319-322.
5. Bailey S. F. and Lacey G. R., 1927, J. Bacteriol., 13:183.
6. Ewing, 1986, Edwards and Ewings Identification of the Enterobacteriaceae, 4th Ed., Elsevier Science Publishing Co., Inc., N.Y.
7. Isenberg, H. D. (ed.). Clinical microbiology procedures handbook, vol.1. American Society for Microbiology, Washington, D.C.
8. Murray, P. R., E. J. Baron, M. A. Pfaller, F. C. Tenover, and R. H. Tenover (eds.). Manual of clinical microbiology, 6th ed. American Society for Microbiology, Washington, D. C.
9. Bacteriological Analytical Manual. 1995. 8th ed. AOAC International, Gaithersburg, M.D.

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITE DM127PSS,QAD/FR/024,Rev.00

Unit 38/39, Kalpataru Industrial Estate,

Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.

Ph: +91-9320126789/9833630009/9819991103

Email: sales@micromasterlab.com

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.