



PRODUCT SPECIFICATION SHEET

Lowenstein Jensen Medium Base / L.J. Medium (DM135)

Intended Use

Lowenstein Jensen Medium Base / L.J. Medium (DM135) is recommended for isolation and cultivation of *Mycobacterium* species.

Product Summary and Explanation

Of the egg-based media, Lowenstein Jensen Medium is most commonly used.⁽¹⁾ L.J. Medium was originally formulated by Lowenstein, containing congo red and malachite green dyes⁽²⁾ Jensen⁽³⁾ modified Lowenstein's medium by altering the citrate and phosphate contents, eliminating the congo red dye and by increasing the malachite green concentration. Gruff^(4, 5) further modified L. J. Medium with the addition of two antimicrobics to increase selectivity. This medium supports the growth of a wide variety of Mycobacteria and can also be used for niacin testing.⁽⁶⁾ Solid media used for isolation and cultivation of Mycobacteria are either egg-based or agar-based. Egg-based media contain whole eggs or egg yolk, potato flour, salts and glycerol and are solidified by inspissation. Mycobacterial infections, particularly tuberculosis, are a worldwide health problem. Almost three million people worldwide die of tuberculosis each year.⁽⁷⁾ Non-tuberculous mycobacteria infections have also increased since 1985⁽⁸⁾ At least 25 species of mycobacteria are associated with human disease and produce usually slowly developing, destructive granulomas that may undergo necrosis with ulceration or cavitation. The use of egg-based media for primary isolation of mycobacteria have two significant advantages. First, egg-based media support a wide variety of mycobacteria. Second, growth of mycobacteria on egg media can be used for niacin testing. Liquification of Lowenstein-Jensen Medium can occur if contaminated with proteolytic organisms. Lowenstein-Jensen Medium is a modification of Lowenstein Medium,⁽⁹⁾ modified by Jensen⁽¹⁰⁾ Jensen modified the medium by alternating the citrate and phosphate contents, eliminating congo red dye, and increasing malachite green concentration.⁽¹¹⁾ Lowenstein-Jensen Medium is commonly used in the clinical laboratory to isolate acid fast organisms from sterile and nonsterile sources.⁽¹²⁾ Hughes and Dixon and Cuthbert⁽¹³⁻¹⁴⁾ reported that the addition of pyruvic acid to egg-based media resulted in improved recovery of tubercle bacilli compared to recovery on egg-based media supplemented only with glycerol. Dixon and Cuthbert recommended using pyruvic acid-egg medium in addition to media supplemented with glycerol for optimum recovery of tubercle bacilli from clinical specimens. Additionally, the medium is available with the addition of 5% sodium chloride. Most rapid growers, the slowly growing *M. triviale* and some strains of *M. flavescens* grow on NaCl containing media. The inability of *M. chelonae* subsp. *Chelonae* to grow helps differentiate it from other members of the *M. fortuitum* complex (e.g., *M. chelonae* subsp. *abscessus*). In the semi-quantitative catalase test, mycobacteria can be differentiated into groups, based upon catalase activity.

Principles of the Procedure

L-Asparagine and Potato Starch are sources of nitrogen and vitamins in Lowenstein-Jensen Medium. Monopotassium Phosphate and Magnesium Sulfate enhance organism growth and act as buffers. Glycerol and the Egg Suspension provide fatty acids and protein required for the metabolism of mycobacteria. The coagulation of the egg albumin during sterilization provides a solid medium for inoculation purposes. Sodium Citrate and Malachite Green are selective agents to prevent growth of most contaminants and allow early growth of mycobacteria. Penicillin and Nalidixic acid in the Gruff's Mycobacterial Supplement (MS049) along with malachite green prevents growth of the majority of contaminants surviving decontamination of the specimen while encouraging earliest possible growth of *Mycobacteria*. RNA in the Gruff's Mycobacterial Supplement (MS049) acts as stimulant and help to increase the isolation rate of *Mycobacteria*. Do not add glycerol to the medium if bovine or other glycerophobic strains are to be cultured. Malachite green serves as an inhibitor and also as pH indicator. Formation of blue zone indicates a decrease in pH by gram-positive contaminants (e.g. *Streptococci*) and yellow zones of dye destruction by gram-negative bacilli. Proteolytic contaminants cause localized or complete digestion of medium. Hardy et al recommended each specimen to be inoculated and incubated in triplicate so as (a) To identify saprophytes at room temperature (25°C) (b) To identify presence or absence of pigmentation by photochromogenes and scotochromogenes at 35°C alternately in light and dark as per the type of organism. Routinely cultivation is carried out aerobically at 35°C.

Formula / Liter

Ingredients	Gms / Liter
L-Asparagine	3.60
Monopotassium phosphate	2.40
Magnesium sulphate	0.24





PRODUCT SPECIFICATION SHEET

Magnesium citrate	0.60
Potato starch, soluble	30.00
Malachite green	0.40
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.

Directions

1. Suspend 37.24 grams of the medium in 600 ml of distilled water containing 12 ml glycerol (for bovine bacteria or other glycerophobic organisms additions of glycerol is not desirable).
2. Heat if necessary to dissolve the medium completely. Sterilize by autoclaving at 15 lbs pressure (121°C) for 15 minutes.
3. Meanwhile prepare 1000ml of whole egg emulsion collected aseptically.
4. Aseptically add and mix egg emulsion base and Gruft Mycobacterial Supplement (MS049) (if desired) gently to obtain uniform mixture.
5. Distribute in sterile screw capped tubes. Arrange tubes in a slanted position. Coagulate and inspissate the medium in an inspissator water bath or autoclave at 85°C for 45 minutes.

Quality Control Specifications

Dehydrated Appearance	Greenish blue to peacock blue homogeneous free flowing powder
Prepared Medium	The mixture of sterile basal medium and whole egg emulsion, when inspissated, coagulates to yield pale bluish green coloured, opaque smooth slants
Reaction of 3.7% solution	--
Gel Strength	Firmness observed due to coagulation of egg emulsion

Expected Cultural Response: Cultural characteristics observed in presence of 5-10% Carbon dioxide, with added egg emulsion base, after an incubation at 35-37°C for 2-4 weeks.

Sr. No.	Organisms	Results to be achieved		
		Growth	Growth with Gruft Supplement (MS049)	Colony Characteristics
1.	<i>Mycobacterium avium</i> ATCC 25291	Luxuriant	Good-luxuriant	Smooth, non-pigmented Colonies
2.	<i>Mycobacterium gordonae</i> ATCC 14470	Luxuriant	Good-luxuriant	Smooth, yellow, Orange colonies
3.	<i>Mycobacterium kansasii</i> ATCC 12478	Luxuriant	Good-luxuriant	Photochromogenic, Smooth to rough
4.	<i>Mycobacterium smegmatis</i> ATCC 14468	Luxuriant	Good-luxuriant	Wrinkled, creamy White colonies
5.	<i>M. tuberculosis</i> H37RV ATCC 25618	Luxuriant	Good-luxuriant	Granular, rough, Warty, dry friable colonies

The organisms listed are the minimum that should be used for quality control testing.





PRODUCT SPECIFICATION SHEET

Test Procedure

1. The test procedures are those recommended by the Centers for Disease Control and Prevention (CDC) for primary isolation from specimens containing mycobacteria. 6 N-Acetyl-L-cysteine-sodium hydroxide (NALC-NaOH) solution is recommended as a gentle but effective digesting and decontaminating agent.
2. These reagents are provided as Mycobacterial Specimen Digestion/Decontamination Kits. For detailed decontamination and culturing instructions, consult an appropriate reference.
3. Following inoculation, keep test containers shielded from light and place them in a suitable system providing an aerobic atmosphere enriched with carbon dioxide.
4. Incubate at $35 \pm 2^\circ\text{C}$. Slanted and bottled media should be incubated in a horizontal plane until the inoculum is absorbed.
5. Tubes and bottles should have screw caps loose for the first 3 weeks to permit the circulation of carbon dioxide for the initiation of growth.
6. Thereafter, to prevent dehydration, tighten caps; loosened briefly once a week. Stand tubes upright if space is a problem.
7. NOTE: Cultures from skin lesions suspected to be *M. Marinum* or *M. ulcerans* should be incubated at $25\text{--}33^\circ\text{C}$ for primary isolation; cultures suspected to contain *M. avium* or *M. Xenopi* exhibit optimum growth at $40\text{--}42^\circ\text{C}$.
8. Incubate a duplicate culture at $35\text{--}37^\circ\text{C}$. For LJ Medium with Iron, specimens must first be isolated in pure culture on an appropriate solid medium.
9. Inoculate LJ Medium with Iron with one drop of a barely turbid suspension of the culture to be tested. For the semi-quantitative catalase test, 1 mL of a 1:1 mixture of 10% polysorbate 80 and 30% hydrogen peroxide is added to each inoculated tube after 2 weeks of incubation.
10. The height of the column of bubbles is recorded after 5 minutes as $<45\text{ mm}$ or $>45\text{ mm}$.

Results

1. Cultures should be read within 5-7 days after inoculation and once a week thereafter for up to 8 weeks.

Record Observations:

- Number of days required for colonies to become macroscopically visible. Rapid growers have mature colonies within 7 days; slow growers require more than 7 days for mature colony forms.
 - Pigment production
2. White, cream to buff - Nonchromogenic (NC)
 3. Lemon, yellow, orange, red-Chromogenic (Ch)
 4. Stained smears may show acid-fast bacilli, which are reported only as "acid-fast bacilli" unless definitive tests are performed.
 5. Bottles may be examined by inverting the bottles on the stage of a dissecting microscope.
 6. Read at $10\text{--}60\times$ with transmitted light. Scan rapidly at $10\text{--}20\times$ for the presence of colonies.
 7. Higher magnification ($30\text{--}60\times$) is helpful in observing colony morphology; i.e., serpentine cord-like colonies.
 8. Examine LJ Medium with Iron for rusty-brown colonies with a tan discoloration in the surrounding medium, indicating uptake of the iron.
 9. The presence or absence of growth in the tube of medium containing 5% NaCl aids in the differentiation of mycobacterial isolates.
 10. The salt tolerance test is positive when numerous colonies appear on the control medium and more than 50 colonies grow on the medium containing 5% NaCl.
 11. Colonies on the control medium, but no visible growth on the test medium after a total of 4 weeks of incubation constitutes a negative test.
 12. In the semi-quantitative catalase test, mycobacteria fall into two groups with *M. tuberculosis* falling into the group producing a column of bubbles less than 45 mm in height.

Storage

Store the sealed bottle containing the dehydrated medium at $10\text{--}30^\circ\text{C}$. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration





PRODUCT SPECIFICATION SHEET

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

