



PRODUCT SPECIFICATION SHEET

Buffered Glucose Broth (DM142BS)

Intended use

Buffered Glucose Broth (DM142BS) is recommended for performing Methyl Red and Voges Proskauer tests in differentiation of coliaerogenes group.

Product Summary and Explanation

In 1915, Clark and Lubs⁽¹⁾ demonstrated that the colon-aerogenes family of bacteria could be divided into two groups based on their action in a peptone and dextrose medium. When tested with pH indicator Methyl red, the "coli" group produced high acidity while the "aerogenes" group produced a less acid reaction. The test to detect high acid end product is known as the Methyl Red (MR) test. This test distinguishes those organisms able to form large amounts of acid from glucose so that the pH falls below 4.4 produce low pH level. Voges and Proskauer⁽²⁾ described a red fluorescent coloration which appeared after the addition of potassium hydroxide to cultures of certain organisms in glucose medium. The red colour produced by the addition of potassium hydroxide to cultures is due to the ability of organisms to produce a neutral product acetoin (acetyl methyl carbinol) from dextrose.⁽³⁾ The acetoin is oxidized in the presence of oxygen and alkali to produce diacetyl which reacts with creatine to produce a red colour and this test was known as VP test. This formulation is also recommended by BIS⁽⁴⁾ and ISO committee⁽⁵⁾ for the detection of coli-aerogenes group. For the detection of *E. coli*, *Vibrio parahaemolyticus* and *Bacillus cereus* responsible for food poisoning a slightly modified formulation (DM142BS) is recommended by BIS.^(6,7,8) To test *V.parahaemolyticus* for VP, addition of 2-3% Sodium chloride to the medium is required.

Principles of the Procedure

Buffered Glucose Broth contains peptic digest of animal tissue which provides carbon, nitrogen, and vitamin used for general growth requirements. Dextrose is the fermentable carbohydrate. Dipotassium phosphate is a buffering agent.

Formula / Liter

Ingredients	Gms / Litre
Peptic digest of animal tissue	5.00
Dextrose	5.00
Dipotassium phosphate	5.00
Final pH: 7.5 ± 0.1 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.

Directions

1. Suspend 15 grams of the medium in 1000 ml of distilled water.
2. Heat if necessary to dissolve the medium completely.
3. Distribute in test tubes in 3 ml amounts or as desired.
4. Autoclave at 121°C, 15 psi pressure, for 15 minutes / validated cycle.

Quality Control Specifications

Dehydrated Appearance	Cream coloured homogeneous free flowing powder
Prepared Medium	Light yellow coloured clear solution without any precipitate





PRODUCT SPECIFICATION SHEET

Reaction of 1.5% Solution	pH : 7.5 ± 0.1 at 25°C
Gel Strength	Not Applicable

Expected Cultural Response: Cultural characteristics observed after an incubation at 30-32°C for 18-48 hours.

Sr. No.	Organisms	Results to be achieved			
		Inoculum (CFU)	Growth	MR Test	VP Test
1.	<i>Bacillus cereus</i> ATCC 10876	50-100	good-luxuriant	negative reaction, yellow colour	positive reaction, eosin pink / red colour within 2-5 minutes
2.	<i>Enterobacter aerogenes</i> ATCC 13048	50-100	good-luxuriant	negative reaction, yellow colour	positive reaction, eosin pink / red colour within 2-5 minutes
3.	<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	positive reaction, bright red colour	negative reaction, no colour change
4.	<i>Klebsiella pneumoniae</i> ATCC 23357	50-100	good-luxuriant	negative reaction, yellow colour	positive reaction, eosin pink / red colour within 2-5 minutes
5.	<i>Salmonella Typhi</i> ATCC 6539	50-100	good-luxuriant	positive reaction, bright red colour	negative reaction, no colour change
6.	<i>Vibrio parahaemolyticus</i> ATCC 17802	50-100	poor	negative reaction, yellow colour	negative reaction, no colour change

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

- Methyl Red (MR) test is performed after maximum of 5 days of incubation at 30°C.⁽⁹⁾
- Voges-Proskauer test (VP) cultures are incubated at 30°C for 24-48 hours.⁽¹⁰⁾
- A variety of other tests have been suggested by Werkman,⁽¹¹⁾ OMeara⁽¹²⁾ Levine, Epstein and Voughn⁽¹³⁾ and Voughn, Mitchell and Levine.⁽⁹⁾
- Werkmans Test : Add 2 drops of a 2% solution of ferric chloride to 50 ml culture and 5 ml of 10% sodium hydroxide. Shake the tube to mix well. Stable copper colour developing in a few minutes is positive reaction.⁽⁵⁾
- OMeara Test : Add of 25 mg of solid creatine to 5ml culture and then add 5 ml concentrated (40%) sodium hydroxide. Red colour development in a few minutes after shaking the tube well, is a positive reaction.⁽⁵⁾
- Levine, Epstein and Voughn⁽¹³⁾ modified OMeara technique by dissolving the creatine in a concentrated solution of potassium hydroxide.
- Voughn, Mitchell and Levine⁽⁹⁾ recommended the method of Barritt⁽¹⁴⁾ as, addition of 1 ml of 40% potassium hydroxide and 3 ml of 5% a - naphthol in absolute ethanol to 5 ml culture. Positive test is indicated by eosine pink colour within 2-5 minutes.
- Refer to appropriate references for standard test procedures.

Results

Refer to appropriate references and test procedures for interpretation of results.

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.





PRODUCT SPECIFICATION SHEET

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. The MR-VP reactions are only part of the tests required to identify organisms.
2. Each laboratory should standardize on the inoculum density, volume of broth and the test container size.
3. MRtest require a minimum incubation of 48 hours before the pH indicator is added.
4. When using Barritt's reagent add α -naphthol first and KOH second; do not reverse this order.
5. Vaughn et al.⁽⁹⁾ warned of false positive VP reactions if the completed tests are left standing for over an hour.

Packaging

Product Name : Buffered Glucose Broth

Product Code : DM142BS

Available Pack sizes : 100gm / 500gm

References

1. Clark and Lubs, 1915, J. Inf. Dis., 17 : 160.
2. Voges and Proskauer, 1898, Zeit, Hyg., 28 : 20. ,,
3. MacFaddin, J.F., 1985, Media for Isolation-Cultivation-Identification-Maintenance of Medical Bacteria, Vol. 1, Williams and Wilkins, Baltimore.
4. Bureau of Indian Standards, IS : 5887 (Part - III) 1976.
5. International Organization for Standardization (ISO), 1993, Draft ISO/DIS 6597.
6. Bureau of Indian Standards, IS : 5887 (Part I) 1976, reaffirmed 1986.
7. Bureau of Indian Standards, IS : 5887 (Part IV) 1976.
8. Bureau of Indian Standards, IS : 5887 (Part - V) 1976, reaffirmed 1986.
9. Vaughn, Mitchell and Levine, 1939, J. Am. Water Works Association, 31:993.
10. Kallas, Chinn and Coulter, 1931, J. Bact., 22 : 125.
11. Werkman, 1930, J. Bact., 20 : 121.
12. O'Meara, 1931, J. Path. Bacteriol., 34 : 401.
13. Levine, Epstein and Vaughn, 1934, Am. J. of Publ. Health, 24 : 505.
14. Barritt, 1936, J. Path. Bacteriol., 42 : 441.

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITED

Unit 38/39, Kalpataru Industrial Estate,
Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.
Ph: +91-22-25895505, 4760, 4681. Cell: 9320126789.

Email: micromaster@micromasterlab.com




sales@micromasterlab.com

DM142BSPSSQAD/FR/024,Rev.00/01.01.2018





PRODUCT SPECIFICATION SHEET

Prepared By	Checked By	Approved By
 01.01.2018	 01.01.2018	 01.01.2018
Microbiologist	Head Quality Control	Head Quality Assurance

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.





PRODUCT SPECIFICATION SHEET

