



PRODUCT SPECIFICATION SHEET

MacConkey Agar W/ 0.15% Bile Salts, CV and NaCl (DM143)

Intended Use

MacConkey agars are slightly selective and differential plating media mainly used for the detection and isolation of gram-negative organisms from clinical, dairy, food, water, pharmaceutical and industrial sources. MacConkey Agar is used for isolating and differentiating lactose fermenting from lactose-non-fermenting gram-negative enteric bacilli. MacConkey Agar Base is used with added carbohydrate in differentiating coliforms based on fermentation reactions. MacConkey Agar without Crystal Violet is used for isolating and differentiating enteric microorganisms while permitting growth of staphylococci and enterococci. The medium can be used also to separate *Mycobacterium fortuitum* and *M. chelonae* from other rapidly growing mycobacteria. MacConkey Agar without Crystal Violet or Salt and MacConkey Agar without Salt are used for isolating and differentiating gram-negative bacilli while suppressing the swarming of most *Proteus* species.

Product Summary and Explanation

MacConkey Agar is based on the bile salt-neutral red-lactose agar of MacConkey. The original MacConkey medium was used to differentiate strains of *Salmonella typhosa* from members of the coliform group. Formula modifications improved the growth of *Shigella* and *Salmonella* strains. These modifications included the addition of 0.5% sodium chloride, decreased agar content, and altered bile salts and neutral red concentrations. The formula improvements gave improved differential reactions between these enteric pathogens and the coliform group. MacConkey Agar contains crystal violet and bile salts that inhibit gram-positive organisms and allow gram-negative organisms to grow. Isolated colonies of coliform bacteria are pink to red in color and may have a zone of precipitated bile. This bile precipitate is due to a local pH drop around the colony due to lactose fermentation. Colonies that do not ferment lactose remain colorless. When lactose non-fermenters grow in proximity to coliform colonies, the surrounding medium appears as cleared areas. It is recommended in the USP for use in the performance of Microbial Limit Tests.

MacConkey Agar Base is prepared without added carbohydrates, which permits their addition either individually or in combination. It is recommended that carbohydrates such as sucrose or lactose be added in a concentration of 1% to the basal medium. MacConkey Agar without Crystal Violet is a differential medium that is less selective than MacConkey Agar. The lack of crystal violet permits the growth of *Staphylococcus* and *Enterococcus*. Staphylococci produce pale pink to red colonies and enterococci produce compact tiny red colonies either on or beneath the surface of the medium. The medium is used also to separate *Mycobacterium fortuitum* and *M. chelonae* from other rapidly growing mycobacteria. MacConkey Agar without Crystal Violet or Salt and MacConkey Agar without Salt (which also lacks crystal violet) are differential media used for isolating and cultivating gram-negative enteric organisms and gram-positive cocci from waters, feces and other sources suspected of containing these organisms, as well as limiting the swarming of *Proteus* species.

Principles of the Procedure

Peptones are sources of nitrogen and other nutrients. Lactose is a fermentable carbohydrate. When lactose is fermented, a local pH drop around the colony causes a color change in the pH indicator (neutral red) and bile precipitation. Bile salts, bile salts no. 3, oxgall and crystal violet are selective agents that inhibit growth of gram-positive organisms. Agar is the solidifying agent.

Formula / Liter

Ingredients	Gms / Litre
Pancreatic digest of gelatin	17.00
Casein enzymic hydrolysate	1.50
Peptic digest of animal tissue	1.50
Lactose	10.00
Bile salts	1.50
Sodium chloride	5.00
Neutral red	0.03
Crystal violet	0.001





PRODUCT SPECIFICATION SHEET

Agar	15.00
Final pH: 7.1 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.

Directions

1. Suspend 51.55 grams of the powder in 1 L of Distilled water. Mix thoroughly.
2. Heat with frequent agitation and boil for 1-2 minutes to completely dissolve the powder.
3. Autoclave at 121°C, 15 psi pressure, for 15 minutes.

Quality Control Specifications

Dehydrated Appearance	Light yellow to pink colored, homogeneous, free flowing powder
Prepared Medium	Red with purplish tinge, clear to slightly opalescent.
Reaction of 5.15% Solution	pH 7.1 ± 0.2 at 25°C
Gel Strength	Firm, compared to 1.5% Agar Gel.

Expected Cultural Response: Cultural response on MacConkey Agar at 35 - 37°C after 18 - 72 hours incubation. The recovery rate is considered as 100% for bacteria growth on Soyabean Casein Digest Agar

Sr. No.	Organisms	Results to be achieved				
		Inoculum (CFU)	Growth	Observed Lot value (CFU)	Recovery	Colour of colony
1.	<i>Escherichia coli</i> ATCC 8739	50-100	Luxuriant	25-100	≥50 %	Pink-red with Bile precipitate
2.	<i>Escherichia coli</i> ATCC 25922	50-100	Luxuriant	25 -100	≥50 %	Pink-red with Bile precipitate
3.	<i>Escherichia coli</i> NCTC 9002	50-100	Luxuriant	25 -100	≥50 %	Pink-red with Bile precipitate
4.	<i>Enterobacter aerogenes</i> ATCC 13048	50-100	Luxuriant	25 -100	≥50 %	Pink to red
5.	<i>Enterococcus faecalis</i> ATCC 29212	50-100	Fair to good	15-40	30 -40 %	Colourless to Pale pink
6.	<i>Salmonella typhimurium</i> ATCC 14028	50-100	Luxuriant	25 -100	≥50 %	Colourless
7.	<i>Staphylococcus aureus</i> ATCC 6538	≥10 ³	Inhibited	0	0%	--
8.	<i>Staphylococcus aureus</i> ATCC 25923	≥10 ³	Inhibited	0	0%	--
9.	<i>Salmonella enteritidis</i> ATCC 13076	50-100	Luxuriant	25 -100	≥50 %	Colourless
10.	<i>Salmonella paratyphi A</i> ATCC 9150	50-100	Luxuriant	25 -100	≥50 %	Colourless
11.	<i>Salmonella paratyphi B</i> ATCC 8759	50-100	Luxuriant	25 -100	≥50 %	Colourless
12.	<i>Salmonella typhi</i> ATCC 6539	50-100	Luxuriant	25 -100	≥50 %	Colourless



PRODUCT SPECIFICATION SHEET

13.	<i>Salmonella abony</i> NCTC 6017	50-100	Luxuriant	25 -100	≥50 %	Colourless
14.	<i>Proteus vulgaris</i> ATCC 13315	50-100	Luxuriant	25 -100	≥50 %	Colourless
15.	<i>Shigella flexneri</i> ATCC 12022	50-100	Fair to good	15-40	30-40%	Colourless
16.	<i>Staphylococcus epidermidis</i> ATCC 12228	≥10 ³	Inhibited	0	0%	--
17.	<i>Corynebacterium diphtheriae</i> type gravis	≥10 ³	Inhibited	0	0%	--

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

For procedures on the isolation and identification of enteric organisms consult the appropriate references.

Results

Lactose-fermenting organisms grow as pink to red colonies with or without a zone of precipitated bile. Lactose-non-fermenting organisms grow as colorless or clear colonies. Swarming by *Proteus* spp. is reduced on MacConkey agars without salt. On MacConkey Agar without Crystal Violet and MacConkey agars without salt, staphylococci produce pale pink to red colonies and enterococci produce tiny red colonies; these organisms are inhibited on MacConkey Agar. On MacConkey Agar without Crystal Violet, potentially pathogenic rapid growers of the *M. fortuitum* complex usually grow in 5-11 days, while the commonly saprophytic species are inhibited. On MacConkey agars without salt, the swarming of *Proteus* is reduced.



MacConkey Agar W/ 0.15% Bile Salts, CV and NaCl (DM143)

Lactose fermenting colonies of *Escherichia coli* ATCC 25922 & Non-lactose fermenting colonies of *Salmonella typhi* ATCC 6539

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature.

Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. Although MacConkey media are selective primarily for gram-negative enteric bacilli, biochemical and serological testing using pure cultures are recommended for complete identification. Appropriate references should be consulted for further information.



PRODUCT SPECIFICATION SHEET

2. Incubation of MacConkey Agar plates under increased CO₂ has been reported to reduce the growth and recovery of a number of strains of gram-negative bacilli.

Packaging

Product Name : MacConkey Agar W/ 0.15% Bile Salts, CV and NaCl

Product Code : DM143

Available Pack sizes : 100gm / 500gm

References

1. Bopp, Brenner, Wells and Strockbine. 1999. In Murray, Baron, Pfaller, Tenover and Tenover (ed.), Manual of clinical microbiology, 7th ed. American Society for Microbiology, Washington, D.C.
2. Flowers, Andrews, Donnelly and Koenig. 1993. In Marshall (ed.), Standard methods for the examination of dairy products. 16th ed., American Public Health Association, Washington, D.C.
3. Downes and Ito (ed.). 2001. Compendium of methods for the microbiological examination of foods, 4th ed. American Public Health Association, Washington, D.C.
4. U.S. Food and Drug Administration. 1995. Bacteriological analytical manual, 8th ed. AOAC International, Gaithersburg, Md.
5. Clesceri, Greenberg and Eaton (ed.). 1998. Standard methods for the examination of water and wastewater, 20th ed. American Public Health Association, Washington, D.C.
6. United States Pharmacopeial Convention, Inc. 2001. The United States pharmacopeia 25/The national formulary 20 - 2002. United States Pharmacopeial Convention, Inc., Rockville, Md.
7. Horwitz (ed.). 2000. Official methods of analysis of AOAC International, 17th ed. AOAC International, Gaithersburg, Md.
8. MacConkey. 1905. J. Hyg. 5:333.
9. Kent and Kubica. 1985. Public health mycobacteriology: a guide for the level III laboratory. USDHHS, Centers for Disease Control, Atlanta, Ga.
10. Master. 1994. In Isenberg (ed.), Clinical microbiology procedures handbook, vol. 1, suppl. 1. American Society for Microbiology, Washington, D.C.
11. Mazura-Reetz, Neblett and Galperin. 1979. Abstr. C179, p. 339. Abstr. Annu. Meet. American Society for Microbiology 1979.

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITED

DM143PSS,QAD/FR/024,Rev.00

Unit 38/39, Kalpataru Industrial Estate,

Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.

Ph: +91-9320126789/9833630009/9819991103

Email: sales@micromasterlab.com

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

