



## PRODUCT SPECIFICATION SHEET

### MacConkey Sorbitol Agar (Sorbitol Agar) (DM154)

#### Intended Use

MacConkey Sorbitol Agar (Sorbitol Agar) (DM154) is recommended for isolation and identification of enteropathogenic *Escherichia coli* strains associated with infant diarrhoea.

#### Product Summary and Explanation

MacConkey Sorbitol Agar (Sorbitol Agar) is based on the formula by Rappaport and Henig.<sup>(1)</sup> This medium is originally developed for isolating enteropathogenic (EPEC) serotypes, O 11 and O55, this medium is also recommended for the isolation and differentiation of enterohemorrhagic *E. coli* O157:H7. This organism causes hemorrhagic colitis, that result from the action of a shiga-like toxin (SLT),<sup>(2)</sup> which results in bloody diarrhea and can lead to kidney failure and death.<sup>(3)</sup> Serotype O157 has been implicated in serious foodborne diseases. On standard MacConkey Agar containing lactose, this strain is indistinguishable from other lactose-fermenting *E. coli*. Unlike most *E. coli* strains, *E. coli* O157:H7 ferments sorbitol slowly or not at all. Therefore, the efficacy of MacConkey Agar containing sorbitol instead of lactose as a differential medium for the detection of *E. coli* O157:H7 in stool cultures was determined. Field trial results showed that the growth of *E. coli* O157:H7 on MacConkey Sorbitol Agar (Sorbitol Agar) was heavy and occurred in almost pure culture as colorless sorbitol-nonfermenting colonies. Most organisms of the fecal flora ferment sorbitol and appear pink on this medium. MacConkey Sorbitol Agar (Sorbitol Agar), therefore, permits ready recognition of *E. coli* O157:H7 in stool cultures.<sup>(3-5)</sup> The addition of cefixime and tellurite significantly reduces the number of sorbitol non-fermenters that need to be screened during the attempted isolation of *E. coli* O157:H7.<sup>(6, 7)</sup>

#### Principles of the Procedure

Peptic digest of animal tissue and proteose peptone in MacConkey Sorbitol Agar, supply necessary nutrients like nitrogenous and carbonaceous compounds, minerals, vitamins and trace ingredients for the growth of organisms. Crystal violet and bile salt mixture present in the medium inhibit growth of gram-positive bacteria, especially enterococci and staphylococci. Sodium chloride maintains osmotic equilibrium. Differentiation of enteric microorganisms is achieved by the combination of D-sorbitol and the neutral red indicator. Colorless or pink to red colonies are produced depending upon the ability of the isolate to ferment the carbohydrate sorbitol.

#### Formula / Liter

Ingredients	Gms / Liter
Peptic digest of animal tissue	17.00
Proteose peptone	3.00
D-Sorbitol	10.00
Bile salts mixture	1.50
Sodium chloride	5.00
Neutral red	0.03
Crystal violet	0.001
Agar	13.50
Final pH: 7.1 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

#### Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.





## PRODUCT SPECIFICATION SHEET

### Directions

1. Suspend 50.03 grams of the medium in one liter of distilled water.
2. Heat to boiling, to dissolve the medium completely.
3. Autoclave at 121°C, 15 psi pressure, for 15 minutes / validated cycle.
4. AVOID OVERHEATING.
5. Cool to 40-50°C and pour into sterile petri plates.

### Quality Control Specifications

Dehydrated Appearance	Light yellow to pink homogeneous free flowing powder
Prepared Medium	Purplish red coloured clear to slightly opalescent gel forms in Petri plates
Reaction of 5.0% Solution	pH : 7.1 ± 0.2 at 25°C
Gel Strength	Firm, comparable with 1.35% Agar gel

**Expected Cultural Response:** Cultural characteristics observed after an incubation 35 - 37°C for 18-24 hours.

Sr. No.	Organisms	Results to be achieved			
		Inoculum (CFU)	Growth	Recovery	Colour of colony
1.	<i>Escherichia coli O157:H7 NCTC 29900</i>	50 -100	good-luxuriant	≥50 %	Colourless
2.	<i>Escherichia coli ATCC 25922</i>	50 -100	good-luxuriant	≥50 %	Pink
3.	<i>Escherichia coli serotype O11 and O55</i>	50-100	fair to good	≥50 %	Colourless
4.	<i>Salmonella Typhi ATCC 6539</i>	50 -100	good-luxuriant	≥50 %	Pink
5.	<i>Shigella flexneri ATCC 12022</i>	50 -100	good-luxuriant	≥50 %	Colourless

The organisms listed are the minimum that should be used for quality control testing.



(Enlarged view)

### MacConkey Sorbitol agar (DM154)

1. *E. coli O157:H7* : smaller, colorless colonies sometimes with an orange halo.
2. *E. coli non-O157:H7* : larger, Pink colonies.

### Test Procedure

Refer to appropriate references for specific procedures, for isolation and identification of enteropathogenic *Escherichia coli* strains associated with infant diarrhoea.

### Results

1. *E. coli O157:H7*, and other organisms that do not ferment sorbitol, are colorless on MacConkey Sorbitol Agar.
2. Sorbitol-fermenting organisms produce pink colonies.
3. Confirmatory biochemical and serological testing should be performed on suspected colonies.





## PRODUCT SPECIFICATION SHEET

---

### Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

### Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

### Limitations of the Procedure

1. Colonies that are sorbitol positive can revert, and can be mistaken for sorbitol negative.
2. Prolonged incubation of the culture may result in colonies of *E. coli* serotype O157:H7 losing their characteristic colorless appearance. There are additional species of facultative anaerobic gram-negative rods that do not ferment sorbitol.

### Packaging

**Product Name : MacConkey Sorbitol Agar (Sorbitol Agar).**

**Product Code : DM154**

**Available Pack sizes : 100gm / 500gm**

### References

1. Rappaport, F., and E. Henig. 1952. Media for the isolation and differentiation of pathogenic *Escherichia coli* (serotypes O111 and O55). *J. Clin. Pathol.* 5:361-362.
2. March and Ratnam. 1986. *J. Clin. Microbiol.* 23:869.
3. U.S. Food and Drug Administration. 1995. Bacteriological analytical manual, 8<sup>th</sup> ed., AOAC International, Gaithersburg, MD.
4. Centers for Disease Control. 1991. Morbid. Mortal. Weekly Rep. 40:265.
5. Bopp, Brenner, Wells and Stockbine. 1999. *In* Murray, Baron, Pfaller, Tenover and Tenover (ed.), Manual of clinical microbiology, 7th ed. American Society for Microbiology, Washington, D.C.
6. Zadik, Chapman and Siddons. 1993. *J. Med. Microbiol.* 39:155.
7. Sanderson, Gay, Hancock, Gay, Fox and Besser. 1995. *J. Clin. Microbiol.* 33:2616.

### Further Information

For further information please contact your local MICROMASTER Representative.



**MICROMASTER LABORATORIES PRIVATE LIMITED**

Unit 38/39, Kalpataru Industrial Estate,  
Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.  
Ph: +91-22-25895505, 4760, 4681. Cell: 9320126789.

Email: [micromaster@micromasterlab.com](mailto:micromaster@micromasterlab.com)  
[sales@micromasterlab.com](mailto:sales@micromasterlab.com)

DM154PSS,QAD/FR/024,Rev.00/01.01.2018





## PRODUCT SPECIFICATION SHEET

Prepared By	Checked By	Approved By
 01.01.2018	 01.01.2018	 01.01.2018
Microbiologist	Head Quality Control	Head Quality Assurance

### Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.





## PRODUCT SPECIFICATION SHEET

---

