

PRODUCT SPECIFICATION SHEET

Malt Agar (DM157)

Intended Use

Malt Agar is recommended for detection and isolation of yeast and mould from dairy products, foods and other material. It is also used for stock culture maintenance of yeast and moulds.

Product Summary and Explanation

Media based on malt extract may be considered as general growth substrates due to their richness and nutrient balance. They are very suitable for the cultivation of fastidious microorganisms. With acidic pH, they are used for the isolation, cultivation and maintenance of yeast and moulds. Malt media for yeasts and moulds have been widely used for many years. In 1919, Reddish⁽¹⁾ prepared a satisfactory substitute for beer wort from malt extract. Thom and Church used Reddish's medium for their studies of *Aspergilli*.⁽²⁾ Malt Agar was employed by Fullmer and Grimes during studies of yeasts on synthetic media.⁽³⁾ Malt Agar is included in Official Methods of Analysis of AOAC International.⁽⁴⁾ It is recommended by APHA⁽⁵⁾ for use in both antibiotic and acidified standard methods for yeast and mould counts in food. This medium is also used for maintaining stock cultures of fungi.

Principles of the Procedure

Malt Agar contains malt extract, which provides carbon, protein and nutrient sources required for the growth of microorganisms. The acidified medium inhibits the growth of bacteria and allows optimal growth of yeasts and moulds.

Formula / Liter

| Ingredients | Gms / Liter |
|--|-------------|
| Malt extract | 30.00 |
| Agar | 15.00 |
| Final pH: 5.5 ± 0.2 at 25°C | |
| Formula may be adjusted and/or supplemented as required to meet performance specifications | |

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.
3. Heating process during rehydration and sterilization should be for shorter period as excessive heat causes partial hydrolysis of the agar, which results in inability to gel properly when cooled. Additional 5 grams of agar may be added if desired.

Directions

1. Suspend 45 grams of the medium in one liter of distilled water.
2. Heat to boiling to dissolve the medium completely.
3. Sterilize by autoclaving at 118°C for 15 minutes.
4. AVOID OVERHEATING, as it will result in a softer and darker agar.

Quality Control Specifications

| | |
|---------------------------|--|
| Dehydrated Appearance | Cream to brownish yellow homogeneous free flowing powder |
| Prepared Medium | Light amber coloured clear to slightly opalescent gel forms in tubes or Petri plates |
| Reaction of 4.5% Solution | pH : 5.5 ± 0.2 at 25°C |
| Gel Strength | Firm, comparable with 1.5% Agar gel |

Expected Cultural Response: Cultural characteristics observed after an incubation at 25 - 30°C for 40 - 48 hours.

| Sr. No. | Organisms | Results to be achieved | | |
|---------|--|------------------------|----------------|----------|
| | | Inoculum (CFU) | Growth | Recovery |
| 1. | <i>Aspergillus brasiliensis</i> ATCC 16404 | 50 - 100 | good-luxuriant | -- |



PRODUCT SPECIFICATION SHEET

| | | | | |
|----|---|----------|----------------|-------|
| 2. | <i>Candida albicans</i> ATCC 10231 | 50 - 100 | good-luxuriant | >=70% |
| 3. | <i>Saccharomyces cerevisiae</i> ATCC 9763 | 50 - 100 | good-luxuriant | >=70% |

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Refer to appropriate references for specific procedures.

Results

Refer to appropriate references and procedures for results.

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. For identification, organisms must be in pure culture. Morphological, biochemical and/or serological tests should be performed for final identification.
2. Consult appropriate texts for detailed information and recommended procedures.

Packaging

Product Name : Malt Agar.

Product Code : DM 157

Available Pack sizes : 500gm

References

1. Reddish, 1919, Abstr. Bacteriol., 3:6.
2. Thom and Church. 1926. The aspergilli. Williams & Wilkins, Baltimore, Md.
3. Fulmer and Grimes. 1923. J. Bacteriol. 8:585.
4. Williams, (Ed.), 2005, Official Methods of Analysis of the Association of Official Analytical Chemists, 19th Ed., AOAC, Washington, D.C.
5. Downes F. P. and Ito K., (Eds.), 2001, Compendium of Methods for the Microbiological Examination of Foods, 4th Ed., APHA, Washington, D.C.

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITED

DM157PSS,QAD/FR/024,Rev.00

Unit 38/39, Kalpataru Industrial Estate,
Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.

Ph: +91-9320126789/9833630009/9819991103

Email: sales@micromasterlab.com

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

