



PRODUCT SPECIFICATION SHEET

SS Agar (Salmonella Shigella Agar) (DM236)

Intended Use

Salmonella Shigella Agar (DM236) is used for the isolation of *Salmonella* spp. and some strains of *Shigella* spp

Product Summary and Explanation

Salmonellosis continues to be an important public health problem worldwide. Infection with non-typhi *Salmonella* often causes a mild, self-limiting illness. Typhoid fever, caused by *Salmonella typhi*, is characterized by fever, headache, diarrhea, abdominal pain, and can result in fatal respiratory, hepatic, and or neurological damage.⁽¹⁾ This infection can result from the consumption of raw, undercooked, or improperly processed foods contaminated with *Salmonella* spp. Shigellosis, caused by *Shigella* spp., is an intestinal illness characterized by abdominal pain, fever, and watery diarrhea. When associated with outbreaks, shigellosis is usually transmitted through contaminated food and/or water.

Salmonella Shigella Agar is essentially a modification of the Desoxycholate Citrate Agar described by Leifson.⁽²⁾ *SS Agar* is moderately selective and differential media for the isolation of pathogenic enteric bacilli, especially those belonging to the genus *Salmonella*. It is not recommended for the primary isolation of *Shigella*. This media has been developed for the selection and differentiation of enteric microorganisms from clinical and non-clinical materials and inhibits the growth of gram-positive species to a varying degree due to the presence of bile salts, brilliant green and sodium citrate.

The high selectivity of *Salmonella Shigella Agar* allows the use of large inocula directly from faeces, rectal swabs or other materials suspected of containing pathogenic enteric bacilli. On fermentation of lactose by few lactose-fermenting normal intestinal flora, acid is produced which is indicated by change of colour from yellow to red by the pH indicator-neutral red. Thus these organisms grow as red pigmented colonies. Lactose non-fermenting organisms grow as translucent colourless colonies with or without black centres. Growth of *Salmonella* species is uninhibited and appears as colourless colonies with black centres resulting from H₂S production. *Shigella* species also grow as colourless colonies which do not produce H₂S. It is recommended to inoculate plates of less inhibitory media parallel to *SS Agar*, such as Hektoen Enteric Agar (DM422) or Deoxycholate Citrate Agar (DM577) for easier isolation of *Shigella* species⁽⁸⁾. *Salmonella Shigella Agar* is recommended for testing clinical specimens and food testing for the presence of *Salmonella* spp. and some *Shigella* spp.^(1,4,5)

Principles of the Procedure

Beef extract and Peptic digest of animal tissue provide sources of nitrogen, carbon, and vitamins required for organism growth. Differentiation of enteric organisms is achieved by the incorporation of lactose in the medium. Organisms that ferment lactose produce acid which, in the presence of the neutral red indicator, results in the formation of red colonies. Lactose non-fermenters form translucent, colorless colonies. The latter group contains the majority of the intestinal pathogens, including *Salmonella* and *Shigella*. Bile Salts, Sodium Citrate and Brilliant Green inhibit Gram-positive bacteria, most coliform bacteria, and inhibit swarming of *Proteus* spp., while allowing *Salmonella* spp. to grow. The sodium thiosulfate and ferric citrate enable the detection of hydrogen sulfide production as evidenced by colonies with black centers. Neutral Red is the pH indicator.

Formula / Liter

Ingredients	Gms / Litre
Beef extract	5.00
Peptic digest of animal tissue	5.00





PRODUCT SPECIFICATION SHEET

Lactose	10.00
Bile salts mixture	8.50
Sodium citrate	10.00
Sodium thiosulphate	8.50
Ferric citrate	1.00
Brilliant green	0.00033
Neutral red	0.025
Agar	15.00
Final pH: 7.0 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. HARMFUL. Harmful if swallowed, inhaled, or absorbed through the skin. May cause allergic reaction and breathing difficulties. Irritating to eyes, skin, and respiratory system.

Directions

1. Suspend 63.02 grams in 1000 ml distilled/purified water.
2. Boil with frequent agitation to dissolve the medium completely.
3. DO NOT AUTOCLAVE OR OVERHEAT.
4. Overheating may destroy selectivity of the medium. Cool to about 50°C.
5. Mix and pour into sterile Petri plates

Quality Control Specifications

Dehydrated Appearance	Light yellow to pink, homogeneous, free flowing powder
Prepared Medium	Reddish-Orange, clear to slightly opalescent gel forms in Petri plates
Reaction of 6.3% Solution	pH 7.0 ± 0.2 at 25°C
Gel Strength	Firm, compared to 1.5% Agar Gel.

Expected Cultural Response: Cultural response on Salmonella Shigella Agar at 35 - 37°C after 18 - 24 hours incubation.

Sr. No.	Organisms	Inoculum (CFU)	Growth	Percentage Recovery	Color of Colonies
1.	<i>Escherichia coli</i> ATCC 25922	50-100	Fair	20-30%	Pink with bile precipitate
2.	<i>Enterobacter aerogenes</i> ATCC 13048	50-100	Fair	20-30%	Cream pink
3.	<i>Enterococcus faecalis</i> ATCC 29212	50-100	None-poor	<=10%	Colourless
4.	<i>Proteus mirabilis</i> ATCC 25933	50-100	Fair-good	30-40%	Colourless
5.	<i>Salmonella choleraesuis</i> ATCC 12011	50-100	Good-luxuriant	>=50%	Colourless, may have black cP entre
6.	<i>Salmonella typhi</i> ATCC 6539	50-100	Good-luxuriant	>=50%	Colourless with black centre






PRODUCT SPECIFICATION SHEET

7.	<i>Salmonella typhimurium</i> ATCC 14028	50-100	Good-luxuriant	>=50%	Colourless with black centre
8.	<i>Salmonella enteritidis</i> ATCC 13076	50-100	Good-luxuriant	>=50%	Colourless with black centre
9.	<i>Shigella flexneri</i> ATCC 12022	50-100	Good	40-50%	Colourless

The organisms listed are the minimum that should be used for quality control testing.



Escherichia coli ATCC 25922 + *Salmonella enteritidis* ATCC 13076

Test Procedure

1. Inoculate the medium heavily with the specimen, spreading a portion of the original inoculum in order to obtain well separated colonies on some part of the plate.
2. For isolation of *Salmonella* spp. and *Shigella* spp. from clinical specimens, inoculate fecal samples and rectal swabs onto one quadrant of Salmonella Shigella Agar, streak for isolation.
3. Incubate plates at 35°C, and examine after 24 and 48 hours for colonies resembling *Salmonella* spp. or *Shigella* spp.
4. For food specimens consult appropriate references for food testing.
5. In parallel with the SS Agar plate, inoculate a tube of Selenite Broth (DM241) enrichment medium, incubate for 12 hours at 35°C, and sub-culture on to another SS Agar plate.
6. A nonselective medium should also be streaked to increase the chance of recovery when the population of gram-negative organisms is low and to provide an indication of other organisms present in the specimen.

Results

1. Enteric organisms are differentiated by their ability to ferment lactose.
2. *Salmonella* spp. and *Shigella* spp. are non-lactose fermenters and form translucent, colorless colonies on Salmonella Shigella Agar. H₂S positive *Salmonella* spp. produce black-center colonies.
3. Some *Shigella* spp. are inhibited on Salmonella Shigella Agar.
4. *E. coli* produces pink to red colonies and may have some bile precipitation





PRODUCT SPECIFICATION SHEET

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. Salmonella Shigella Agar is highly selective and not recommended as the primary isolation of Shigella.^(1,2,6) Some Shigella spp. May be inhibited.
2. A few nonpathogenic organisms may grow on Salmonella Shigella Agar. These organisms can be differentiated by their ability to ferment lactose and other confirmatory tests.
3. This medium is highly selective and resistant strains of shigellae will not grow on it. It is not recommended for the primary isolation of shigellae^(2,6)
4. Media recommended for the isolation of *Shigella* are Hektoen Enteric and XLD Agars⁽⁸⁾

Packaging

Product Name: SS agar (Salmonella Shigella Agar)

Product Code : DM236

Available Pack sizes : 100gm / 500gm

References

1. P. R. Murray, E. J. Baron, M. A. Pfaller, F. C. Tenover, and R. H. Tenover (eds.). 1995. Manual of clinical microbiology, 6th ed. American Society for Microbiology, Washington, D. C.
2. Leifson, E. 1935. New culture media based on sodium desoxycholate for the isolation of intestinal pathogens and for the enumeration of colon bacilli in milk and water. J. Pathol. Bacteriol 40:581.
3. Rose, H. M., and M. H. Kolodny. 1942. The use of SS (Shigella-Salmonella) Agar for the isolation of Flexner Dysentery bacilli from the feces. J. Lab. Clin. Med. 27:1081-1083.
4. Isenberg, H. D. (ed.). 1992. Interpretation of aerobic bacterial growth on primary culture media, Clinical microbiology procedures handbook, vol. 1 p. 1.61-1.67. American Society for Microbiology, Washington, D.C.
5. Vanderzant, C., and D. F. Splittstoesser (eds.). 1992. Compendium of methods for the microbiological examination of foods, 3rd ed. American Public Health Association, Washington, D.C.
6. Taylor, W. I., and B. Harris. 1965. Isolation of shigellae. II. Comparison of plating media and enrichment broths. Am. J. Clin. Pathol. 44:476.
7. McFaddin, J. F. 1985. Media for isolation-cultivation-identification-maintenance of medical bacteria, Vol. 1. Williams & Wilkins, Baltimore, MD.
8. Pollock and Dahlgren. 1974. Appl. Microbiol. 27:197.

Further Information

For further information please contact your local MICROMASTER Representative.








PRODUCT SPECIFICATION SHEET



MICROMASTER LABORATORIES PRIVATE LIMITED
Unit 38/39, Kalpataru Industrial Estate,
Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.
Ph: +91-22-25895505, 4760, 4681. Cell: 9320126789.
Email: micromaster@micromasterlab.com
sales@micromasterlab.com

DM236PSS,QAD/FR/024,Rev.00/01.01.2018

Prepared By	Checked By	Approved By
 01.01.2018	 01.01.2018	 01.01.2018
Microbiologist	Head Quality Control	Head Quality Assurance

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

