



## PRODUCT SPECIFICATION SHEET

### Xylose-Lysine Deoxycholate Agar (DM297I)

#### Intended Use

Xylose-Lysine Deoxycholate Agar (DM297I) is recommended for selective isolation and enumeration of *Salmonella typhi* and other *Salmonella* species from pharmaceutical products using microbial limit testing in compliance with IP.

#### Product Summary and Explanation

The Enterobacteriaceae is a large family of Gram-negative bacteria that includes, along with many harmless symbionts, many of the more familiar pathogens, such as *Salmonella*, *Escherichia coli*, *Yersinia pestis*, *Klebsiella* and *Shigella*. Other disease-causing bacteria in this family include *Proteus*, *Enterobacter*, *Serratia*, and *Citrobacter*. They normally inhabit the intestines of humans and animals.<sup>(1)</sup> Human *Salmonella* infections are most commonly caused by ingestion of food, water or milk, contaminated by human or animal excreta.<sup>(2)</sup> A wide variety of media have been developed to aid in the selective isolation and differentiation of enteric pathogens. Due to the large numbers of different microbial species and strains with varying nutritional requirements and chemical resistance patterns, investigators have developed various formulae to meet general as well as specific needs relative to isolation and identification of the microorganisms. Xylose Lysine Deoxycholate Agar is a selective as well as differential medium formulated by Taylor<sup>(3-7)</sup> for the isolation and identification of enteric pathogens especially Shigellae from stool samples. This medium is also employed for pharmaceutical testing and non-sterile product testing for the detection (or absence) of *Salmonella* after enrichment in Rappaport Vassiliadis Salmonella Enrichment Broth (DM1405I). This medium is prepared according to IP<sup>(8)</sup> and is in accordance with the harmonized method of USP/BP/EP.<sup>(9-11)</sup>

#### Principles of the Procedure

Xylose-Lysine Deoxycholate Agar contains yeast extract which is a source of nitrogen, carbon, and vitamins required for organism growth. Deoxycholate, ferric ammonium citrate and sodium thiosulphate are selective agents that inhibit gram-positive microorganisms. Xylose, lactose, and sucrose are fermentable carbohydrate sources. Almost all the enteric bacteria except Shigellae, ferment xylose fermented which enables the differentiation of Shigellae from Salmonellae. Salmonellae metabolize the xylose and decarboxylate lysine and thus changing the pH to alkaline and mimic Shigellae reaction. Lactose and sucrose are added in excess to produce acid and hence non-pathogenic H<sub>2</sub>S producers do not decarboxylate lysine thereby preventing Shigellae reaction by lysine positive coliforms. Sodium thiosulphate prevents the desiccation of these compounds during storage by reactivating sulphur containing compounds. It also forms the substrate for enzyme thiosulphate reductase, which breaks it to form H<sub>2</sub>S. Thiosulphate and ferric ammonium citrate forms H<sub>2</sub>S indicator system. Sodium thiosulphate is also inactivator of halogens, mercurial and aldehyde and can minimize its toxicity in the testing sample, if any during microbial limit tests. Sodium chloride maintains the osmotic equilibrium in this medium. Phenol red is the pH indicator.

#### Formula / Liter

Ingredients	Gms / Liter
Xylose	3.50
L-Lysine	5.00
Lactose monohydrate	7.50
Sucrose	7.50
Sodium chloride	5.00
Yeast extract	3.00
Phenol red	0.08
Sodium deoxycholate	2.50
Sodium thiosulphate	6.80
Ferric ammonium citrate	0.80
Agar	13.50
Final pH: 7.4 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	



## PRODUCT SPECIFICATION SHEET

### Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.

### Directions

1. Suspend 54.8 g (the equivalent weight of dehydrated medium per litre) in one liter of purified water.
2. Heat with frequent agitation until the medium boils.
3. DO NOT OVERHEAT OR AUTOCLAVE.
4. Transfer immediately to a water bath at 50°C. After cooling, pour into sterile Petri plates.
5. It is advisable not to prepare large volumes, which will require prolonged heating and may produce precipitate.

### Quality Control Specifications

Dehydrated Appearance	Light yellow to pink homogeneous free flowing powder
Prepared Medium	Red coloured clear to slightly opalescent gel forms in Petri plates
Reaction of % Solution	Not Applicable
Gel Strength	Firm, comparable with 1.35 % Agar gel

### Cultural Response

Growth Promotion is carried out in accordance with the harmonized method of IP. Cultural response was observed after an incubation at 30-35°C for specified time. Recovery rate is considered as 100% for bacteria growth on Soyabean Casein Digest Agar.

### Growth promoting properties

Growth of microorganism comparable to that previously obtained with previously tested and approved lot of medium occurs at the specified temperature for not more than the shortest period of time specified inoculating  $\leq 100$  cfu(at 30-35°C for  $\leq 18$  hours).

### Indicative properties

Colonies are comparable in appearance and indication reaction to those previously obtained with previously tested and approved lot of medium occurs for the specified temperature for a period of time within the range specified inoculating  $\leq 100$ cfu (at 30-35°C for 18-72 hours).

### Inhibitory properties

No growth of the test microorganism occurs for the specified temp for not less than longest period of time specified inoculating  $\geq 100$ cfu (at 30-35°C for  $\geq 72$  hours).

**Expected Cultural Response:** Cultural characteristics observed after incubation at 30-35 °C for 18-48 hours. Recovery rate is considered as 100% for bacteria growth on Soyabean Casein Digest Agar.

Sr. No.	Organisms	Results to be achieved					
		Inoculum (CFU)	Growth	Observed Lot value(CFU)	Recovery	Colour of Colony	Incubation period
	<b>Growth Promoting + Indicative</b>						
1.	<i>Salmonella Typhimurium</i> ATCC 14028	50 -100	luxuriant	25 -100	$\geq 50$ %	red with black centres	18 -72 hrs
2.	<i>Salmonella Abony</i> NCTC 6017	50 -100	good-luxuriant	25 -100	$\geq 50$ %	red with black centres	18 -72 hrs
	<b>Additional Microbiological testing</b>						



## PRODUCT SPECIFICATION SHEET

3.	<i>Escherichia coli</i> ATCC 8739	50-100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
4.	<i>Escherichia coli</i> ATCC 25922	50-100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
Sr. No.	Organisms	Results to be achieved					
		Inoculum (CFU)	Growth	Observed Lot value(CFU)	Recovery	Colour of Colony	Incubation period
5.	<i>Escherichia coli</i> NCTC 9002	50-100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
6.	<i>Proteus vulgaris</i> ATCC 13315	50 -100	good-luxuriant	25 -100	≥50 %	grey with black centres	18 -72 hrs
7.	<i>Salmonella Paratyphi A</i> ATCC 9150	50 -100	good-luxuriant	25 -100	≥50 %	red	18 -72 hrs
8.	<i>Salmonella Paratyphi B</i> ATCC 8759	50 -100	good-luxuriant	25 -100	≥50 %	red with black centers	18 -72 hrs
9.	<i>Salmonella Enteritidis</i> ATCC 13076	50 -100	good-luxuriant	25 -100	≥50 %	red with black centers	18 -72 hrs
10.	<i>Salmonella Typhi</i> ATCC 6539	50 -100	good-luxuriant	25 -100	≥50 %	red with black centers	18 -72 hrs
11.	<i>Shigella dysenteriae</i> ATCC 13313	50 -100	good-luxuriant	25 -100	≥50 %	red	18 -72 hrs
12.	<i>Shigella flexneri</i> ATCC 12002	50 -100	fair-good	15 -40	30 -40 %	red	18 -72 hrs
13.	<i>Shigella sonnei</i> ATCC 25931	50 -100	fair-good	15 -40	30 -40 %	red	18 -72 hrs
14.	<i>Enterobacter aerogenes</i> ATCC 13048	50 -100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
15.	<i>Enterobacter cloacae</i> ATCC 13047	50 -100	fair	10 -30	20 -30 %	yellow	18 -72 hrs
16.	<i>Staphylococcus aureus</i> ATCC 25923	≥10 <sup>3</sup>	inhibited	0	0%	--	≥72 hrs
17.	<i>Staphylococcus aureus</i> ATCC 6538	≥10 <sup>3</sup>	inhibited	0	0%	--	≥72 hrs
18.	<i>Enterococcus faecalis</i> ATCC 29212	≥10 <sup>3</sup>	inhibited	0	0%	--	≥72 hrs

The organisms listed are the minimum that should be used for quality control testing.

### Test Procedure

Refer to appropriate references for standard test procedures.

### Results

- Degradation of fermentable sugars proceed concurrently and generates acids, which cause pH indicator to give various shades of colour, causing a colour change in the colonies and in the medium from red to yellow on prolonged incubation.
- Hydrogen sulfide production results in colonies with black centers under alkaline conditions, which can be inhibited by acid production by carbohydrate fermentation. Alkaline condition causes the colour of the medium to change back to red.
- This medium is an ideal medium for screening samples containing mixed flora of enteric pathogens as recovery of Salmonellae and Shigellae is not conspicuous by even profuse growth of other species.<sup>(12,13)</sup>



## PRODUCT SPECIFICATION SHEET

---

### Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

### Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

### Limitations of the Procedure

1. Red, false-positive colonies may occur with some *Proteus* and *Pseudomonas* species.
2. Incubation in excess of 48 hours may lead to false-positive results.
3. *S. Paratyphi A*, *S. Choleraesuis*, *S. pullorum* and *S. gallinarum* may form red colonies without black centers, thus resembling *Shigella* species.
4. Some *Proteus* strains will give black-centered colonies on XLD Agar.
5. Consult appropriate texts for detailed information and recommended procedures.

### Packaging

**Product Name :** Xylose-Lysine Deoxycholate Agar

**Product Code :** DM297I

**Available Pack sizes :** 100gm / 500gm

### References

1. Pelczar M. J. Jr., Reid R. D., Chan E. C. S., 1977, Microbiology, 4th Ed., Tata McGraw-Hill Publishing Company Ltd, New Delhi.
2. Koneman E. W., Allen S. D., Janda W. M., Schreckenberger P. C., Winn W. C. Jr., 1992, Colour Atlas and Textbook of Diagnostic Microbiology, 4th Ed., J. B. Lippincott Company.
3. Taylor W. L., 1965, Am. J. Clin. Pathol., 44:471-475.
4. Taylor W. L. and Harris B., 1965, Am. J. Clin. Pathol., 44:476.
5. Taylor W. L. and Harris B., 1967, Am. J. Clin. Pathol., 48:350.
6. Taylor W. L. and Schelhart B., 1967, Am. J. Clin. Pathol., 48:356.
7. Taylor W. L. and Schelhart B., 1968, Am. J. Clin. Pathol., 16:1387.2.
8. Indian Pharmacopoeia, 2010 Ministry of Health and Family Welfare, Govt. of India.
9. The United States Pharmacopoeia, 2011, The United States Pharmacopoeial Convention. Rockville, MD.
10. British Pharmacopoeia, 2011, The Stationery office British Pharmacopoeia.
11. European Pharmacopoeia, 2011, European Dept. for the quality of Medicines.
12. McCarthy M.D., 1966, N.Z. J. Med. Lab. Technol., 20:127.
13. Isenberg H.D., Kominos S. and Siegal M., 1969, Appl. Microbiol., 18:656.

### Further Information

For further information please contact your local MICROMASTER Representative.



**MICROMASTER LABORATORIES PRIVATE LIMITED**

DM297IPSS,QAD/FR/024,Rev.00/01.01.2018

Unit 38/39, Kalpataru Industrial Estate,  
Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.  
Ph: +91-22-25895505, 4760, 4681. Cell: 9320126789.

Email: [micromaster@micromasterlab.com](mailto:micromaster@micromasterlab.com)  
[sales@micromasterlab.com](mailto:sales@micromasterlab.com)

---



## PRODUCT SPECIFICATION SHEET

Prepared By	Checked By	Approved By
 01.01.2018	 01.01.2018	 01.01.2018
Microbiologist	Head Quality Control	Head Quality Assurance

### Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.