

PRODUCT SPECIFICATION SHEET

King's OF Medium Base (DM331)

Intended Use

King's OF Medium Base (DM331) is recommended for studying oxidation-fermentation of carbohydrates by Campylobacter species.

Product Summary and Explanation

Campylobacter is a gram-negative, motile bacterium that cause a gastrointestinal infection called campylobacteriosis when it gets lodged in the walls of intestine. The bacteria normally inhabit the intestinal tract of warm-blooded animals such as poultry and cattle, and are frequently detected in foods derived from these animals. Although raw milk is a frequently reported vehicle of outbreaks of Campylobacter enteritis, studies have revealed that mishandled poultry is more important than raw milk in transmitting Campylobacter jejuni enteritis. (1,2,3) For the identification of an organism genus and species the utilization pattern for several carbohydrates (e.g. lactose, maltose, xylose, sucrose etc) is often needed. Kings OF Medium is designed as recommended by APHA for studying the oxidation-fermentation reaction of carbohydrates by Campylobacter species. (4)

Principles of the Procedure

Carbon Utilisation Agar contains casein enzymic hydrolysate which is a source of nitrogenous compounds required for the growth of *Campylobacter* species. Phenol red is the pH indicator. Oxidation of carbohydrate is indicated by a yellow colour formation.

Formula / Liter

Ingredients	Gms / Liter	
Casein enzymic hydrolysate	0.20	
Phenol red	0.003	
Agar	0.30	
Final pH: 7.4 ± 0.2 at 25°C		
Formula may be adjusted and/or supplemented as required to meet performance specifications		

Precautions

- 1. For Laboratory Use only.
- 2. IRRITANT. Irritating to eyes, respiratory system, and skin.

Directions

- 1. Suspend 0.5 grams of the medium in one litre of distilled water.
- 2. Heat if necessary to dissolve the medium completely.
- 3. Autoclave at 121°C, 15 psi pressure, for 15 minutes / validated cycle.
- 4. Cool to $40-50^{\circ}C$ and aseptically add filter sterilized solution of desired carbohydrate to get a final concentration of 1% and dispense in sterile tubes.

Quality Control Specifications

Dehydrated Appearance Light yellow to beige homogeneous free flowing powder	
Prepared Medium	Light pink coloured, clear to slightly opalescent gel forms in tubes as butts
Reaction of 0.05% Solution pH : 7.4 ± 0.2 at $25^{\circ}C$	
Gel Strength	Semisolid,comparable with 0.03% Agar gel









PRODUCT SPECIFICATION SHEET

Expected Cultural Response: Cultural characteristics observed with added Dextrose under reduced oxygen atmosphere, after an incubation at 42°C for 24-48 hours.

Sr	Onconismo	Results to be achieved	
No	Organisms	Growth	Acid (with dextrose)
1.	Campylobacter jejuni ATCC 29428	good-luxuriant	positive reaction, yellow colour

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Refer appropriate references for specific test procedures.

Results

The medium will be yellow (acid) when removed from the microaerobic atmosphere due to CO_2 absorption. To read OF reactions, let the tubes stand at room temperature until the OF control becomes neutral or alkaline, usually within 2 hours. Refer appropriate references and test procedures for interpretation of results.

Storage

Store the sealed bottle containing the dehydrated medium at 2 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

- 1. For identification, organisms must be in pure culture. Morphological, biochemical and/or serological tests should be performed for final identification.
- 2. Consult appropriate texts for detailed information and recommended procedures.

Packaging

Product Name : King's OF Medium Base

Product Code: DM331

Available Pack sizes: 100gm / 500gm

References

- Shirling E. B. and Gottlieb D., 1966, Methods for Characterization of Streptomyces species, Int. J. Syst. Bacteriol., 16:313.
- 2. Atlas R. M., 1993, Handbook of Microbiological Media, Parks L.C. (Ed.), CRC Press, Inc.

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITED

DM331PSS,QAD/FR/024,Rev.00

Unit 38/39, Kalpataru Industrial Estate,

Off G.B. Road, Near 'R-Mall', Thane (W) - 400607. M.S. INDIA.

Ph: +91-9320126789/9833630009/9819991103.

Email: sales@micromasterlab.com









PRODUCT SPECIFICATION SHEET

Disclaimer:

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.





