



PRODUCT SPECIFICATION SHEET

M17 Broth (DM565)

Intended Use

M17 Broth (DM565) is recommended for cultivating and enumerating *lactic streptococci* in yogurt, cheese starters and other dairy products.

Product Summary and Explanation

Lactic streptococci are acid-producing bacteria and are nutritionally fastidious, requiring complex culture media for optimum growth. One study showed that in a synthetic medium, all strains had an obligate requirement for at least six amino acids and three vitamins.⁽¹⁾ These homofermentative lactic streptococci produce large amounts of acid and, in a culture medium without an adequate buffering system, the pH decreases and adversely affects growth. M16 Medium was developed by Lowrie and Pearce⁽²⁾ but it lacked a strong buffering system. Terzaghi and Sandine⁽³⁾ worked with M16 Medium and demonstrated that the rapid drop in pH that accompanies lactic streptococcal growth can adversely affect colony size and phage plaque formation. They modified M16 Medium using disodium-β-glycerophosphate as a buffer and called it M17. Disodium glycerophosphate maintains the pH above 5.7. The maintenance of pH is very important as the lower pH results in injury and reduced recovery of lactic Streptococci. Glycerophosphate does not form precipitate with calcium which is needed for the plaque assay of lactic bacteriophages. Shankar and Davies⁽⁴⁾ found that disodium-β-glycerophosphate in M17 Broth suppressed *Lactobacillus bulgaricus* and selectively isolated *Streptococcus thermophilus* from yogurt.⁽⁵⁾

Principles of the Procedure

M17 Broth contains peptic digest of animal tissue, casein enzymic hydrolysate, papaic digest of soyabean meal, yeast extract, beef extract, provide carbonaceous, nitrogenous compounds, vitamin B complex and other essential growth factors. Lactose is the fermentable carbohydrate and ascorbic acid is stimulatory for the growth of lactic Streptococci. Magnesium sulphate provides essential ions to the organisms. Disodium-3-glycerophosphate maintains the pH above 5.7.

Formula / Liter

Ingredients	Gms / Liter
Peptic digest of animal tissue	2.50
Casein enzymic hydrolysate	2.50
Papaic digest of soyabean meal	5.00
Yeast extract	2.50
Beef extract	5.00
Lactose	5.00
Ascorbic acid	0.50
Disodium - β - glycerophosphate	19.00
Magnesium sulphate	0.25
Final pH: 7.1 ± 0.1 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.





PRODUCT SPECIFICATION SHEET

Directions

1. Suspend 42.25 grams of the medium in one liter of distilled water.
2. Heat if necessary to dissolve the medium completely.
3. Autoclave at 121°C, 15 psi pressure, for 15 minutes / validated cycle.
4. Mix well and dispense as desired.

Quality Control Specifications

Dehydrated Appearance	Cream to yellow homogeneous free flowing powder
Prepared Medium	Light yellow coloured clear to slightly opalescent solution in tubes
Reaction of 4.23% Solution	pH : 7.1 ± 0.1 at 25°C
Gel Strength	Not Applicable

Expected Cultural Response: Cultural characteristics observed after an incubation at 35-37°C for 24-48 hours.

Sr. No.	Organisms	Results to be achieved	
		Inoculum (CFU)	Growth
1.	<i>Enterococcus faecalis</i> ATCC 29212	50-100	good-luxuriant
2.	<i>Lactobacillus bulgaricus</i> ATCC 11842	50-100	none-poor
3.	<i>Lactobacillus leichmannii</i> ATCC 4797	50-100	good-luxuriant
4.	<i>Lactobacillus plantarum</i> ATCC 8014	50-100	good-luxuriant
5.	<i>Streptococcus thermophilus</i> ATCC 14485	50-100	good-luxuriant

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Refer to appropriate references for standard test procedures.

Results

Refer to appropriate references and test procedures for interpretation of results.

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. For identification, organisms must be in pure culture. Morphological, biochemical and/or serological tests should be performed for final identification.
2. Consult appropriate texts for detailed information and recommended procedures.





PRODUCT SPECIFICATION SHEET

Packaging

Product Name : M17 Broth

Product Code : DM565

Available Pack sizes : 100gm/ 500gm

References

1. Reiter and Oram. 1962. J. Dairy Res. 29:63.
2. Lowrie and Pearce. 1971. J. Dairy Sci. Technol. 6:166.
3. Terzaghi and Sandine. 1975. Appl. Microbiol. 29:807.
4. Shankar and Davies. 1977. J. Soc. Dairy Tech. 30:28.
5. International Dairy Federation. 1981. Identification and enumeration of microorganisms in fermented milks. Joint IDF/ISO/AOAC Group E44.

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITED


DM565PSS, QAD/FR/024, Rev.00/01.01.2018

Unit 38/39, Kalpataru Industrial Estate,
Near Runwal Estate, Behind 'R-Mall' , Ghodbunder Road,
Thane (W) - 400607. M.S. INDIA.

Ph: +91-22-25895505, 4760, Cell: 9320126789.

Email: micromaster@micromasterlab.com

sales@micromasterlab.com

Prepared By	Checked By	Approved By
 01.01.2018	 01.01.2018	 01.01.2018
Microbiologist	Head Quality Control	Head Quality Assurance

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

