



PRODUCT SPECIFICATION SHEET

Enterobacteria Enrichment Broth, Mossel (DM616E)

Intended Use

Enterobacteria Enrichment Broth, Mossel (DM616E) is recommended for selective enrichment of *Enterobacteriaceae* from pharmaceutical products using the microbial limit testing in compliance with EP.

Product Summary and Explanation

The enumeration of *Enterobacteriaceae* is of great concern in monitoring the sanitary condition of food. The *Enterobacteriaceae* family consists of *Salmonella*, *Shigella* and other enteric pathogens. These organisms find entry into the food system through faecally contaminated water. *Enterobacteriaceae* can be injured in food-processing procedures, which include exposure to low temperatures, sub-marginal heat, drying, radiation, preservatives or sanitizers. Recovery relies on proper resuscitation of damaged cells.⁽¹⁾ The resuscitation procedure is recommended for dried foods,⁽²⁾ animal feeds⁽³⁾ and semi-preserved foods.⁽⁴⁾ Mossel et al⁽⁵⁾ formulated EE Broth, Mossel, which is recommended as an enrichment medium for bile tolerant gram-negative bacteria in the biological examination of foods, animal feed stuffs.^(5,6) This medium is prepared as per European Pharmacopoeia⁽⁹⁾ and is in accordance with the harmonized method of USP/BP/EP/JP/IP.^(7,8,9,10,11)

Principles of the Procedure

Enterobacteria Enrichment Broth, Mossel contains pancreatic digest of gelatin and glucose monohydrate which provides nitrogen, vitamins and amino acids for the growth of members of *Enterobacteriaceae*. Brilliant green and ox bile, purified are selective agents which inhibit the growth of gram-positive bacteria. Phosphates form the buffering system of the medium and neutralizes acids produced by lactose fermenters that otherwise would adversely affect the growth of the organism. Lactose is replaced by glucose in this medium as lactose negative, anaerogenic lactose-positive or late lactose fermenting *Enterobacteriaceae* are often missed by the standard Coli-aerogenes test. The cells damaged while drying or low pH is resuscitated in well-aerated Soybean Casein Digest Broth (DM277E) for 2 hours at 25°C prior to enrichment in EE Broth.

Formula / Liter

Ingredients	Gms / Liter
Pancreatic digest of gelatin	10.00
Glucose monohydrate	5.00
Dehydrated ox-bile	20.00
Disodium hydrogen phosphate, dihydrate	8.00
Potassium dihydrogen phosphate	3.00
Brilliant green	0.015
Final pH: 7.2 ± 0.2 at 25°C	
Formula may be adjusted and/or supplemented as required to meet performance specifications	

Precautions

1. For Laboratory Use only.
2. IRRITANT. Irritating to eyes, respiratory system, and skin.

Directions

1. Suspend 42.93 grams of the medium in one liter of purified/distilled water.
2. Dispense 120 ml amounts in 250 ml flasks or 9 ml amounts in tubes.
3. Stopper with cotton plugs or loose fitting caps.
4. Heat in free flowing steam or boiling water for 30 minutes.
5. Avoid overheating of the medium. DO NOT AUTOCLAVE.





PRODUCT SPECIFICATION SHEET

Quality Control Specifications

Dehydrated Appearance	Light yellow to greenish yellow homogeneous free flowing powder
Prepared Medium	Emerald green coloured, clear solution without any precipitate
Reaction of 4.29% Solution	pH 7.2 ± 0.2 at 25°C
Gel Strength	Not Applicable

Growth Promotion Test

Growth Promotion is carried out in accordance with the harmonized method of EP. Cultural response was observed after an incubation at 30-35°C for specified time.

Growth promoting properties

Clearly visible growth of microorganism comparable to that previously obtained with previously tested and approved lot of medium occurs at the specified temperature for not more than the shortest period of time specified inoculating ≤100 cfu (at 30-35°C for ≤24 hours).

Inhibitory properties

No growth of the test microorganism occurs for the specified temp for not less than longest period of time specified inoculating ≥ 100cfu (at 30-35°C for ≥ 48 hours).

Expected Cultural Response: Cultural response was observed after an incubation at 30-35°C for 24-48 hours.

Sr. No.	Organisms	Results to be achieved			
		Inoculum (CFU)	Growth	Acid	Incubation Temperature /period
Growth Promoting					
1.	<i>Escherichia coli</i> ATCC 8739	50 -100	good-luxuriant	positive reaction, yellow colour	30 -35 °C ≤24 hrs
2.	<i>Pseudomonas aeruginosa</i> ATCC 9027	50 -100	good-luxuriant	positive reaction, yellow colour	30 -35 °C ≤24 hrs
Inhibitory					
3.	<i>Staphylococcus aureus</i> ATCC 6538	≥10 ³	inhibited	--	30 -35 °C ≥48 hrs
Test for Enterobacteriaceae					
4.	<i>Escherichia coli</i> ATCC 8739	50-100	Luxuriant	positive reaction, yellow colour	30 -35 °C 18-48hrs
5.	<i>Pseudomonas aeruginosa</i> ATCC 25783	50-100	Luxuriant	positive reaction, yellow colour	30 -35 °C 18-48hrs
Additional Microbiological Testing					
6.	<i>Escherichia coli</i> ATCC 25922	50-100	good-luxuriant	positive reaction, yellow colour	30 -35 °C 24 -48 hrs
7.	<i>Escherichia coli</i> NCTC 9002	50 -100	good-luxuriant	positive reaction, yellow colour	30 -35 °C 24 -48 hrs
8.	<i>Pseudomonas aeruginosa</i> ATCC 27853	50 -100	good-luxuriant	positive reaction, yellow colour	30 -35 °C 24 -48 hrs
9.	<i>Enterobacter aerogenes</i> ATCC 14028	50 -100	good-luxuriant	positive reaction, yellow colour	30 -35 °C 24 -48 hrs
10.	<i>Proteus mirabilis</i>	50 -100	good-luxuriant	positive reaction,	30 -35 °C





PRODUCT SPECIFICATION SHEET

	<i>ATCC 25933</i>			yellow colour	24 -48 hrs
11.	<i>Salmonella Enteritidis</i> <i>ATCC 13076</i>	50 -100	good-luxuriant	positive reaction, yellow colour	30 -35 °C 24 -48 hrs
12.	<i>Shigella boydii ATCC 12030</i>	50 -100	good-luxuriant	negative reaction no colour change	30 -35 °C 24 -48 hrs
13.	<i>Staphylococcus aureus</i> <i>ATCC 25923</i>	$\geq 10^3$	inhibited	--	30 -35 °C ≥ 48 hrs

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

1. EE Broth is an enrichment broth and should be used in conjunction with Violet Red Bile Glucose Agar (DM287E). A loopful of the enriched sample from EE Broth is sub-cultured onto Violet Red Bile Glucose Agar (DM287E) after an initial incubation at 30-35°C for 24 hours.
2. Typical pink colonies from DM287E are sub-cultured for biochemical confirmation by oxidase and fermentation reactions.⁽⁷⁾ Decimal dilutions of the food homogenate are used if the expected counts are high or else initial suspension is used. EE Broth, Mossel (DM616E)
3. Refer to appropriate references for standard test procedures.

Results

1. A positive reaction results in acid production which causes the colour of EE Broth Mossel to become yellow.
2. A negative reaction results in no colour change and the medium remains green.

Storage

Store the sealed bottle containing the dehydrated medium at 10 - 30°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

1. Some strains may grow poorly or fail to grow on this medium.
2. For identification, organisms must be in pure culture. Morphological, biochemical and/or serological tests should be performed for final identification.
3. Consult appropriate texts for detailed information and recommended procedures.

Packaging

Product Name : Enterobacteria Enrichment Broth, Mossel

Product Code : DM616E

Available Pack sizes : 100gm / 500gm

References

1. Mossel D. A. A., and Harrewijn G. A., 1972, *Alimenta* II, 29-30
2. Mossel D.A.A. and Ratto M.A., 1970, *Appl. Microbiol.*, 20:273.
3. Mossel D.A.A. and Shennan J.L. and Clare V., 1973, *J. Sci. Fd. Agric.*, 24 : 499.
4. Mossel D.A.A., Ratto M.A., 1973, *J. Fd. Technol.*, 8 : 97.
5. Mossel D. A. A., Vissar M. and Cornellsen A. M. R., 1963, *J. Appl. Bacteriol.*, 26(3):444.
6. Van Schothurst M. et al, 1966, *Vet Med.*, 13(3):273.
7. The United States Pharmacopoeia, 2011, The United States Pharmacopoeial Convention. Rockville, MD.
8. British Pharmacopoeia, 2011, The Stationery office British Pharmacopoeia
9. European Pharmacopoeia, 2011, European Dept. for the quality of Medicines.
10. Japanese Pharmacopoeia, 2008.
11. The Indian Pharmacopoeia 2010, Govt. of India, 2010. The Controller of Publication, Delhi.





PRODUCT SPECIFICATION SHEET

Further Information

For further information please contact your local MICROMASTER Representative.




MICROMASTER LABORATORIES PRIVATE LIMITED

DM616EPSS, QAD/FR/024, Rev.00/01.01.2018

Unit 38/39, Kalpataru Industrial Estate,
Near Runwal Estate, Behind 'R-Mall', Ghodbunder Road,
Thane (W) - 400607. M.S. INDIA.

Ph: +91-22-25895505, 4760, Cell: 9320126789.

Email: micromaster@micromasterlab.com
sales@micromasterlab.com

Prepared By	Checked By	Approved By
 01.01.2018	 01.01.2018	 01.01.2018
Microbiologist	Head Quality Control	Head Quality Assurance

Disclaimer :

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.

