

PRODUCT SPECIFICATION SHEET

Chocolate Agar Plate (RP003)

Intended Use

For isolation of Neisseria gonorrhoeae from chronic and acute gonococcal infections.

Product Summary and Explanation

The cultivation medium for gonococci should ideally be a rich nutrients base with blood, either partially lysed or completely lysed. Neisseria gonorrhoeae is a gram-negative bacteria and the causative agent of gonorrhoea, however it is also occasionally found in the throat. The diagnosis and control of gonorrhoea have been greatly facilitated by improved laboratory methods for detecting, isolating and studying N. gonorrhoeae. Chocolate Agar Base, with the addition of supplements, gives excellent growth of the gonococcus without overgrowth by contaminating organisms. G.C. Agar (DM116) can also be used in place of Chocolate Agar Base, which gives slightly better results than Chocolate Agar. (1)

Principles of the Procedure

Chocolate Agar Base with addition of supplement not only supports the growth of the gonococcus in pure culture but also permits its development from the mixed flora encountered in chronic gonococcal infections. Carpenter reported that this medium and Haemoglobin (MS088) is useful for cultural detection of the gonococcus.

Ingredients	Gms / Liter			
Proteose peptone	20.00			
Dextrose	0.500			
Sodium chloride	5.000			
Disodium phosphate	5.000			
Haemoglobin	2.00			
Agar	15.00			
Final pH: 7.3 ± 0.2 at 25°C				
Formula may be adjusted and/or supplemented as required to meet performance specifications				

Precautions

- 1. For Laboratory Use only.
- 2. IRRITANT. Irritating to eyes, respiratory system, and skin.

Directions

Either streak, inoculate or surface spread the test inoculum (50-100 CFU) aseptically on the plate.

Quality Control Specifications

Appearance	Sterile Chocolate Agar in 90mm disposable plate		
Color of Final medium	Chocolate Brown colored medium		
Clarity of Final Medium	Opaque gel		
pH at 25°C	7.10 ± 7.50		
Gel Strength	Firm, comparable with 1.7% Agar gel		
Quantity of Medium in Plates	18-22ml		
Sterilization of Basal Medium	Autoclaving at 15 lbs Pressure		
Pre Incubation Check	cubation Check No growth should be observed after incubation at 30-35 35 - 37 $^{\circ}$ C/ for 18 - 24		
	hrs		









PRODUCT SPECIFICATION SHEET

Expected Cultural Response: Cultural characteristics observed after an incubation at 35-37°C for 24 - 48 hours.

	Organism	Incubation time/ temp	Inoculum (<i>C</i> FU)	Growth	Recovery	Remarks
1.	Neisseria gonorrhoeae ATCC 19424	35 - 37 °C/ 24 - 72 hrs.	50 -100	Luxuriant	>=70 %	Complies
2.	Neisseria meningitidis ATCC 13090	35 - 37 °C/ 24 - 72 hrs.	50 -100	Luxuriant	>=70 %	Complies
	Organism	Incubation time/ temp	Inoculum (<i>C</i> FU)	Growth	Recovery	Remarks
3.	Streptococcus pneumonia ATCC 6303	35 - 37 °C/ 24 - 72 hrs.	50-100	Luxuriant	>=70 %	Complies
4.	Streptococcus pyogenes ATCC 19615	35 - 37 °C/ 24 - 72 hrs.	50-100	Luxuriant	>=70 %	Complies
5.	Haemophilus influenza ATCC 19418	35 - 37 °C/ 24 - 72 hrs.	50-100	Luxuriant	»=70 %	Complies

The organisms listed are the minimum that should be used for quality control testing.

Test Procedure

Refer appropriate references for detailed instructions of specific procedures.

Results

Refer to appropriate references and standard procedures for interpretation of results.

Storage

Store the sealed bottle containing the dehydrated medium at 2 - 8°C. Once opened and recapped, place container in a low humidity environment at the same storage temperature. Protect from moisture and light.

Expiration

Refer to the expiration date stamped on the container. The dehydrated medium should be discarded if not free flowing, or if the appearance has changed from the original color. Expiry applies to medium in its intact container when stored as directed.

Limitations of the Procedure

For identification, organisms must be in pure culture, Morphological, biochemical and/ serological tests should be performed for final identification.

Consult appropriate texts for detailed information and recommended procedure.

Packaging

Product Name: Chocolate Agar Plate

Product Code : RP003 Available Pack sizes : 10 PL

References

1. Carpenter C. M., Bucca M. A., Buck T. C., Casman E. P., Vhristensen C. W., Crowe E., Drew R., Hill J., Lankford L. E., Morton H. E., Peizer L. R., Shaw C. J., and Thayer J. D., 1949, Am. J. Syphil. Gonorrh. Veneral Diseases, 33:164. 2. Muench. Wochschr., 80:846:1933.

3.McLeod J. W., Cootes J. C., Happold F. C., Priestely D. P., Wheatley B., 1934, J. Path. Bacteriol., 39:221.

4.J.Infectious Diseases, 61:129:1937.

5.Am. J. Syphillis, 20:347:1936.

6.Am. J. Syphillis, 22:55:1938.









PRODUCT SPECIFICATION SHEET

7.Seventh Annual Year book (1936-37) P.133, supple., Am. J. Pub.Health.27: no.3: 1937. 8.Bull. Genitoinfectious diseases, Mass. State Health Dept., 2:1:1938.

Further Information

For further information please contact your local MICROMASTER Representative.



MICROMASTER LABORATORIES PRIVATE LIMITED

RP003PSS, QAD/FR/024, Rev. 00

Unit 38/39, Kalpataru Industrial Estate,
Off G.R. Dood, Near 'P-Mall', Thong (W) - 400607, M.S. TND:

Off G.B. Road, Near 'R-Mall' , Thane (W) - 400607. M.S. INDIA. Ph: 91-9320126789/9833630009/9819991103

Email: sales@micromasterlab.com

Disclaimer:

All Products conform exclusively to the information contained in this and other related Micromaster Publications. Users must ensure that the product(s) is appropriate for their application, prior to use. The information published in this publication is based on research and development work carried out in our laboratory and is to the best of our knowledge true and accurate. Micromaster Laboratories Pvt Ltd reserves the right to make changes to specifications and information related to the products at any time. Products are intended for laboratory, diagnostic, research or further manufacturing use only and not for human or animal or therapeutic use, unless otherwise specified. Statements included herein should not be considered as a warranty of any kind, expressed or implied, and no liability is accepted for infringement of any patents.





